

विध्न विचारत भीरु जन, नहीं आरम्भे काम, विपति देख छोड़े तुरंत मध्यम मन कर श्याम।
पुरुष सिंह संकल्प कर, सहते विपति अनेक, 'बना' न छोड़े ध्येय को, रघुबर राखे टेक।।

*रचितः मानव धर्म प्रणेता
सद्गुरु श्री रणछोड़दासजी महाराज*

Subject : CHEMISTRY

Available Online : www.MathsBySuhag.com

PHYSICAL CHEMISTRY

Teko[®]
CLASSES
....the support

**Wishing You & Your Family A Very
Happy & Prosperous Deepawali**



**DEEPAWALI
ASSIGNMENT**



Address : Plot No. 27, III- Floor, Near Patidar Studio,
Above Bond Classes, Zone-2, M.P. NAGAR, Bhopal
☎ : 0 903 903 7779, 98930 58881, WhatsApp 9009 260 559
www.TekoClasses.com www.MathsBySuhag.com

SPARKLES

- Q.1 NO_2 is involved in the formation of smog and acid rain. A reaction that is important in the formation of NO_2 is $\text{O}_3(\text{g}) + \text{NO}(\text{g}) \rightleftharpoons \text{O}_2(\text{g}) + \text{NO}_2(\text{g})$ $K_c = 6 \times 10^{34}$, if the air over Bansal Classes contained $1 \times 10^{-6} \text{M O}_3$, $1 \times 10^{-5} \text{M NO}$, $2.5 \times 10^{-4} \text{M NO}_2$ and $8.2 \times 10^{-3} \text{M O}_2$, what can we conclude?
 (A) There will be a tendency to form more NO and O_3
 (B) There will be a tendency to form more NO_2 and O_2
 (C) There will be a tendency to form more NO_2 and O_3
 (D) There will be no tendency for change because the reaction is at equilibrium.
- Q.2 The standard enthalpy of formation of gaseous H_2O at 298 K is -242 kJ mol^{-1} . Calculate ΔH° at 373 K given the following values of the molar heat capacities at constant pressure
 Molar heat capacity of $\text{H}_2\text{O}(\text{g}) = 33.5 \text{ JK}^{-1}\text{mol}^{-1}$
 Molar heat capacity of $\text{H}_2(\text{g}) = 28.8 \text{ JK}^{-1}\text{mol}^{-1}$
 Molar heat capacity of $\text{O}_2(\text{g}) = 29.4 \text{ JK}^{-1}\text{mol}^{-1}$
 Assume that the heat capacities are independent of temperature.
 (A) 508 kJmol^{-1} (B) -242 kJmol^{-1} (C) $-242.75 \text{ kJmol}^{-1}$ (D) None
- Q.3 A reaction takes place in three steps. The rate constant are K_1 , K_2 & K_3 . The overall rate constant $K = \frac{K_1 \times K_3}{K_2}$. If energy of activation are 20, 15 and 10 kJ/mole, the overall energy of activation is
 (A) 10 (B) 15 (C) 20 (D) $\frac{40}{3}$
- Q.4 At 200°C PCl_5 dissociates as follows:
 $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$
 It was found that the equilibrium vapours are 62 times as heavy as hydrogen. The degree of dissociation of PCl_5 at 200°C is:
 (A) 10% (B) 42% (C) 50% (D) 68%
- Q.5 For the reaction
 $\text{C}_2\text{H}_6(\text{g}) \rightleftharpoons \text{C}_2\text{H}_4(\text{g}) + \text{H}_2(\text{g})$
 K_p is 0.05 at 900 K. If an initial mixture comprising 20 mol of C_2H_6 and 80 mol of inert gas is passed over a dehydrogenation catalyst at 900 K, what is the equilibrium mole percentage of C_2H_6 in the gas mixture? The total pressure is kept of 0.5 bar.
 (A) 4.3 (B) 9.67 (C) 8.76 (D) 72.5
- Q.6 A certain radio isotope ${}^A_Z\text{X}$ ($t_{1/2} = 10$ days) decays to give ${}^{A-4}_{Z-2}\text{Y}$, if two mole of ${}^A_Z\text{X}$ is kept in a sealed vessel, how much He will accumulate in 20 days at STP?
 (A) 16.8 litre (B) 22.4 litre (C) 33.6 litre (D) None
- Q.7 For the transformation
 $\text{H}_2\text{O}(l, 1 \text{ atm}) \rightarrow \text{H}_2\text{O}(g, 0.1 \text{ atm})$; $\Delta H_{\text{vap}} = 40.668 \text{ kJ mol}^{-1}$
 The change in entropy ($\text{Jk}^{-1} \text{ mol}^{-1}$) is
 (A) 19.14 (B) 89.89 (C) 109.03 (D) 128.17
- Q.8 When sulphur in the form of S_8 is heated at 900 K, the initial pressure of 1 atm falls by 30% at equilibrium. This is because of conversion of some S_8 to S_2 . The value of equilibrium constant for this reaction is
 (A) 2.96 (B) 2.05 (C) 0.39 (D) 3.9
- Q.9 A solution of 0.4 g sample of H_2O_2 reacted with 0.632 g of KMnO_4 in presence of sulphuric acid. The percentage purity of the sample of H_2O_2 is
 (A) 95% (B) 85% (C) 80% (D) none of these

Q.10 For the reaction (1) and (2)




Given $K_{P_1} : K_{P_2} = 9 : 1$

If the degree of dissociation of A and D be same then the total pressure at equilibrium (1) and (2) are in the ratio.

- (A) 3 : 1 (B) 36 : 1 (C) 1 : 1 (D) 0.5 : 1

Q.11 A 0.518 g sample of lime stone is dissolved in HCl and then the calcium is precipitated as CaC_2O_4 . After filtering and washing the precipitate, it requires 40.0 mL of 0.250 N KMnO_4 solution acidified with H_2SO_4 to titrate it as,



The percentage of CaO in the sample is:

- (A) 54.0% (B) 27.1% (C) 42% (D) 84%

Q.12 The time of decay for a nuclear reaction is given by $t = 4t_{1/2}$. The relation between the mean life (T) and time of decay (t) is given by

- (A) $2T \ln 2$ (B) $4T \ln 2$ (C) $2T^4 \ln 2$ (D) $\frac{1}{T^2} \ln 2$

Q.13 0.3 g of an oxalate salt was dissolved in 100 mL solution. The solution required 90 mL of N/20 KMnO_4 for complete oxidation. The % of oxalate ion in salt is:

- (A) 33% (B) 66% (C) 70% (D) 40%

Q.14 For the reaction




the values of ΔH° and ΔS° at 298 K are 77.2 kJ mol^{-1} and $122 \text{ JK}^{-1} \text{ mol}^{-1}$ respectively. The standard equilibrium constant at the same temperature is

- (A) 0.695×10^{-8} (B) 6.95×10^{-8} (C) 69.5×10^{-8} (D) 695×10^{-8}

Q.15 An element A in a compound AB has oxidation number A^{n-} . It is oxidised by $\text{Cr}_2\text{O}_7^{2-}$ in acidic medium. In the experiment 1.68×10^{-3} mole of $\text{K}_2\text{Cr}_2\text{O}_7$ were used for 3.26×10^{-3} mole of AB. The new oxidation number of A after oxidation is:

- (A) 3 (B) $3 - n$ (C) $n - 3$ (D) $+n$

Q.16 Ethylene is produced by




Cyclobutane

The rate constant is $2.48 \times 10^{-4} \text{ sec}^{-1}$. In what time will the molar ratio of the ethylene to cyclobutane in reaction mixture attain the value 1?

- (A) 27.25 minute (B) 28.25 minute (C) 25 minute (D) 20 minute

Q.17 Rate of formation of SO_3 in the following reaction



is 100 kg min^{-1} . Hence rate of disappearance of SO_2 will be:

- (A) 100 kg min^{-1} (B) 80 kg min^{-1} (C) 64 kg min^{-1} (D) 32 kg min^{-1}

Get Solution of These Packages & Learn by Video Tutorials on www.MathsBySuhag.com


- Q.18 The hydrolysis of an ester was carried out separately with 0.05 N HCl and 0.05 N H₂SO₄. Which of the following will be true?
 (A) $k_{\text{HCl}} > k_{\text{H}_2\text{SO}_4}$ (B) $k_{\text{HCl}} < k_{\text{H}_2\text{SO}_4}$ (C) $k_{\text{HCl}} = k_{\text{H}_2\text{SO}_4}$ (D) $k_{\text{H}_2\text{SO}_4} = 2k_{\text{HCl}}$
- Q.19 The reaction $\text{N}_2 + \text{O}_2 \rightleftharpoons 2\text{NO}$, follows the kinetics Rate $\propto [\text{N}_2][\text{O}_2]$. The order of reactions when both N₂ and O₂ are in small amount, N₂ is in large excess as compared to O₂ and both N₂ and O₂ are in large excess are
 (A) 2, 1 and 2 (B) 2, 1 and 1 (C) 2, 1 and 0 (D) 1, 1 and 0
- Q.20 The reaction: $2\text{NO} + 2\text{H}_2 \longrightarrow \text{N}_2 + 2\text{H}_2\text{O}$ has been assigned to follow following mechanism
 I. $\text{NO} + \text{NO} \rightleftharpoons \text{N}_2\text{O}_2$ (fast)
 II. $\text{N}_2\text{O}_2 + \text{H}_2 \longrightarrow \text{N}_2\text{O} + \text{H}_2\text{O}$ (slow)
 III. $\text{N}_2\text{O} + \text{H}_2 \longrightarrow \text{N}_2 + \text{H}_2\text{O}$ (fast)
 The rate constant of step II is $1.2 \times 10^{-4} \text{ mole}^{-1}\text{L min}^{-1}$ while equilibrium constant of step I is 1.4×10^{-2} . What is the rate of reaction when concentration of NO and H₂ each is 0.5 mole L⁻¹.
 (A) $2.1 \times 10^{-7} \text{ mol L}^{-1} \text{ min}^{-1}$ (B) $3.2 \times 10^{-6} \text{ mol L}^{-1} \text{ min}^{-1}$
 (C) $3.5 \times 10^{-4} \text{ mol L}^{-1} \text{ min}^{-1}$ (D) none of these
- Q.21 In a reaction carried out at 500 K, 0.001% of the total number of collisions are effective. The energy of activation of the reaction is approx.
 (A) 15.8 K cal mol⁻¹ (B) 11.5 K cal mol⁻¹ (C) 12.8 K cal mol⁻¹ (D) zero
- Q.22 The catalyst decrease the E_a from 100 kJ mol⁻¹ to 80 kJ mol⁻¹. At what temperature the rate of reaction in the absence of catalyst at 500 K will be equal to rate reaction in presence of catalyst.
 (A) 400 K (B) 200 K (C) 625 K (D) none of these
- Q.23 The thermal decomposition of acetaldehyde: $\text{CH}_3\text{CHO} \longrightarrow \text{CH}_4 + \text{CO}$, has rate constant of $1.8 \times 10^{-3} \text{ mol}^{-1/2} \text{ L}^{1/2} \text{ min}^{-1}$ at a given temperature. How would $-\frac{d[\text{CH}_3\text{CHOO}]}{dt}$ will change if concentration of acetaldehyde is doubled keeping the temperature constant?
 (A) will increase by 2.828 times (B) will increase by 11.313 times
 (C) will not change (D) will increase by 4 times
- Q.24 If an aqueous solution at 25 °C has twice as many OH⁻ as pure water its pOH will be
 (A) 6.699 (B) 7.307 (C) 7 (D) 6.98
- Q.25 A radioactive mixture containing a short lived species A and short lived species B. Both emitting α-particles at a given instant, emits at rate 10,000 α-particles per minute. 10 minutes later, it emits at the rate of 7000 particles per minute. If half lives of the species are 10 min and 100 hours respectively, then the ratio of activities of A : B in the initial mixture was
 (A) 3 : 7 (B) 4 : 6 (C) 6 : 4 (D) none
- Q.26 In order to determine the volume of blood in an animal without killing it, a 1.00 ml sample of an aqueous solution containing tritium is injected into the animal blood stream. The sample injected has an activity of 1.8×10^6 cps (counts per second). After sufficient time for the sample to be completely mixed with the animal blood due to normal blood circulation, 2.00 ml of blood are withdrawn from animal and the activity of the blood sample withdrawn is found to be 1.2×10^4 cps. Calculate the volume of the animal blood.
 (A) 300 ml (B) 200 ml (C) 250 ml (D) 400 ml

Get Solution of These Packages & Learn by Video Tutorials on www.MathsBySuhag.com

Q.27 A radioactive isotope is being produced at a constant rate x . Half-life of the radioactive substance is y . After sometimes number of radioactive nuclei becomes constant, the value of this constant is

- (A) $\frac{xy}{\ln 2}$ (B) xy (C) $(\ln 2)xy$ (D) x/y

Q.28 A vessel contains $H_2(g)$ at 2 atm pressure, when $H_2S(g)$ at a pressure of 4 atm is introduced into the vessel. Where reaction




Occurs at a temperature of 1000 K. It is found that

$$\left[\frac{n(H_2)}{n(H_2S)} \right]_{\text{at equilibrium}} = \left[\frac{n(H_2S)}{n(H_2)} \right]_{\text{at } t=0}, \text{ then}$$

- (A) maximum weight of solid formed is 32 gm
 (B) maximum weight of solid formed is 0.32 gm
 (C) $K_p = K_c RT$
 (D) $K_c = 256$

Q.29 At a certain temperature the following equilibrium is established



One mole of each of the four gases is mixed in one litre container and the reaction is allowed to reach equilibrium state. When excess of baryta water is added to the equilibrium mixture, the weight of white precipitate obtained is 236.4 gm. The equilibrium constant K_c of the following reaction is

- (A) 1.2 (B) 2.25 (C) 2.1 (D) 3.6

Q.30 Rate of disappearance of the reactant A at two different temperatures is given by A \rightleftharpoons B

$$\frac{-d[A]}{dt} = 2 \times 10^{-2} \text{ sec}^{-1} [A] - 4 \times 10^{-3} \text{ sec}^{-1} [B] \text{ at } 300 \text{ K}$$

$$\frac{-d[A]}{dt} = 4 \times 10^{-2} \text{ sec}^{-1} [A] - 16 \times 10^{-4} \text{ sec}^{-1} [B] \text{ at } 400 \text{ K}$$

heat of reaction in the given temperature range, when equilibrium is set up is

- (A) $\frac{2.303 \times 2 \times 300 \times 400}{100} \log 50 \text{ Cal}$ (B) $\frac{2.303 \times 2 \times 300 \times 400}{100} \log 250 \text{ Cal}$
 (C) $\frac{2.303 \times 2 \times 300 \times 400}{100} \log 5 \text{ Cal}$ (D) None

BOMB

- Q.1 For the reaction,
- $$2\text{SO}_3(\text{g}) \rightleftharpoons 2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \quad \Delta H_1^\circ = \text{standard enthalpy change of reaction}$$
- $$\Delta S_1^\circ = \text{standard entropy at temperature T}$$
- $$2\text{NO}_3(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g}) + \text{O}_2(\text{g}) \quad \Delta H_2^\circ = \text{standard enthalpy at temperature T}$$
- $$\Delta S_2^\circ = \text{standard entropy at temperature T}$$


Find out the equilibrium constant for the reaction

$\text{SO}_2(\text{g}) + \text{NO}_3(\text{g}) \rightleftharpoons \text{SO}_3(\text{g}) + \text{NO}_2(\text{g})$ at the same temperature in terms of above parameter.

- Q.2 Given the sub shells 1s, 2s, 2p, 3p and 3d. Identify those meet the following descriptions
- Has l (azimuthal quantum no.) = 2
 - Can have m (magnetic quantum no.) = -1
 - Is empty in a nitrogen atom
 - Is full in a carbon atom
 - Can contain two electrons, both with spin $m_s = +1/2$
 - Contains the outermost shell electrons in oxygen atom



- Q.3 Rate of diffusion of ozonized oxygen is $0.4\sqrt{5}$ times that of pure oxygen. Find
- Percentage by mass of ozone in the ozonized sample
 - Degree of dissociation of oxygen assuming pure O_2 in the sample initially.

- Q.4 A 250 ml flask and 100 ml flask are separated by a stop cock. At 350 K, the nitric oxide in the larger flask exerts a pressure of 0.46 atm and the smaller one contains oxygen at 0.86 atm. The gases are mixed by opening the stop cock. The reaction occurring are




The first reaction is complete while the second one is at equilibrium. Assuming all the gases to behave ideally, calculate the K_p for the equilibrium reaction if the final total pressure is 0.37 atm.

- Q.5 25 ml of a dilute aqueous solution of p-hydroxy benzoic acid is titrated with NaOH (aq); the solution has pH = 4.57, when 8.12 ml of 0.0200 M NaOH had been added and pH = 7.02 after 16.24 ml had been added (the equivalent point). Use these data to determine K_{a1} and K_{a2} for p-hydroxy benzoic acid.



- Q.6 A balloon containing air at 27°C and 1 atm initially is filled with air further isothermally and reversibly till the pressure is 4 atm. If the initial diameter of the balloon is 1 m and the pressure at each state is proportional to diameter of the balloon, calculate the work done.

- Q.7 (a) Thermal decomposition of dibromosuccinic acid (DBSA) taking place according to the following equation, obeys first order kinetics



The progress of reaction may be followed by means of alkali titration of the solution (reaction mixture) at various time intervals. If T_0 and T_t be the titres at zero – time and at any time t , respectively, a and $a - x$ be the concentrations DBSA at zero time and at any time t , respectively, show that


$$\frac{T_0}{3T_0 - 2T_t} = \frac{a}{a - x}$$

- (b) If the rate of a reaction doubles from 0°C to 10°C by what factor would the ratio increase when the temperature increases from 100°C to 110°C and what is the activation energy?

Successful People Replace the words like; "wish", "try" & "should" with "I Will". Ineffective People don't.

Get Solution of These Packages & Learn by Video Tutorials on www.MathsBySuhag.com

Q.8 Compressibility factor Z vs P plot for four real gases A, B, C & D are shown at 300 K




- Out of gases A and B whose molecules are larger? Justify.
- If slope of curve for B is 0.02 atm^{-1} , calculate Van der Waals constant b for gas B.
- Arrange Boyle temperatures (T) for gases A, B, C & D in increasing order.

Q.9 A 5.0 gm sample containing Pb_3O_4 , PbO_2 and some inert impurity is dissolved in 250 ml dil. HNO_3 solution and 2.7 g of $\text{Na}_2\text{C}_2\text{O}_4$ was added so that all lead converted into Pb^{2+} . A 10 ml portion of this solution required 8.0 ml, 0.02 M KMnO_4 for titration of excess of oxalate. In another experiment, 25 ml of solution was taken and excess oxalate was removed by extraction, this required 10 ml of a permanganate solution for oxidation of Pb^{2+} to Pb^{4+} . 10 ml of this permanganate solution is equivalent to 4.48 ml, 5V H_2O_2 solution. Calculate mass % of PbO_2 & Pb_3O_4 in the original sample. [At. wt. of Pb = 207]

Q.10 The apparatus shown consists of three temperature jacketed 1 litre bulbs connected by stopcocks. Bulb A contains a mixture of $\text{H}_2\text{O}(\text{g})$, $\text{CO}_2(\text{g})$ and $\text{N}_2(\text{g})$ at 27°C and a total pressure of 547.2 mm Hg. Bulb B is empty and is held at a temperature -23°C . Bulb C is also empty and is held at a temperature of -173°C . The stopcocks are closed and the volumes of lines connecting the bulbs is zero.

Given: $\text{CO}_2(\text{g})$ converted into $\text{CO}_2(\text{s})$ at -78° , $\text{N}_2(\text{g})$ converted into $\text{N}_2(\text{s})$ at -196°C & $\text{H}_2\text{O}(\text{g})$ converted into $\text{H}_2\text{O}(\text{s})$ at 0°C .

[Use $R = 0.08 \text{ atm-litre/mole}\cdot\text{K}$]



- The stopcock between A & B is opened and the system is allowed to come to equilibrium. The pressure in A & B is now 228 mmHg. What do bulbs A & B contain?
- How many moles of H_2O are in system?
- Both stopcocks are opened and the system is again allowed to equilibrium. The pressure throughout the system is 45.6 mmHg. What do bulbs A, B and C contain?
- How many moles of N_2 are in the system?

ANSWER KEY

SPARKLES

Q.1	B	Q.2	C	Q.3	B	Q.4	D	Q.5	C	Q.6	C	Q.7	D
Q.8	A	Q.9	B	Q.10	B	Q.11	A	Q.12	B	Q.13	B	Q.14	B
Q.15	B	Q.16	A	Q.17	B	Q.18	A	Q.19	A	Q.20	A	Q.21	B
Q.22	A	Q.23	A	Q.24	A	Q.25	C	Q.26	A	Q.27	A	Q.28	D
Q.29	B	Q.30	C										

BOMB

- Q.1
$$K = \exp - \left\{ \frac{(\Delta H_2^\circ - \Delta H_1^\circ) - T(\Delta S_2^\circ - \Delta S_1^\circ)}{2RT} \right\}$$
- Q.2 (a) 3d (b) 2p, 3p, 3d (c) 3s, 3p and 3d (d) 1s & 2s
 (e) 2p, 3p and 3d (f) 2p, 2s
- Q.3 60%, 0.6 Q.4 0.64 atm Q.5 $K_{a1} = 2.7 \times 10^{-5}$, $K_{a2} = 3.4 \times 10^{-10}$
- Q.6 1.015×10^4 kJ Q.7 (b) $k_2/k_1 = 1.454$, 10.64 kcal
- Q.8 (a) $A > B$, (b) 0.492 L/mol, (c) $T_A < T_B < T_C < T_D$
- Q.9 % $Pb_3O_4 = 68.5$, % $PbO_2 = 23.9$
- Q.10 (a) Bulb A : $N_2(g), CO_2(g)$
 Bulb B : $N_2(g), CO_2(g), H_2O(s)$
 (b) No. of mole of $H_2 = 0.0025$
 (c) Bulb A : $N_2(g)$
 Bulb B : $N_2(g), H_2O(s)$
 Bulb C : $N_2(g), CO_2(s)$
 (d) Total no. of mole of N_2 in system = 0.013