

$$a^3 - b^3 = (a - b)(a - \omega b)(a - \omega^2 b) ; \quad x^2 + x + 1 = (x - \omega)(x - \omega^2) ;$$

$$a^3 + b^3 = (a + b)(a + \omega b)(a + \omega^2 b) ;$$

$$a^3 + b^3 + c^3 - 3abc = (a + b + c)(a + \omega b + \omega^2 c)(a + \omega^2 b + \omega c)$$

nth ROOTS OF UNITY :

If $1, \alpha_1, \alpha_2, \alpha_3, \dots, \alpha_{n-1}$ are the n, n^{th} root of unity then :

(i) They are in G.P. with common ratio $e^{i(2\pi/n)}$ &

(ii) $1^p + \alpha_1^p + \alpha_2^p + \dots + \alpha_{n-1}^p = 0$ if p is not an integral multiple of n
 $= n$ if p is an integral multiple of n

(iii) $(1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_{n-1}) = n$ &

$(1 + \alpha_1)(1 + \alpha_2) \dots (1 + \alpha_{n-1}) = 0$ if n is even and 1 if n is odd.

(iv) $1 \cdot \alpha_1 \cdot \alpha_2 \cdot \alpha_3 \dots \alpha_{n-1} = 1$ or -1 according as n is odd or even.

THE SUM OF THE FOLLOWING SERIES SHOULD BE REMEMBERED :

$$(i) \cos \theta + \cos 2\theta + \cos 3\theta + \dots + \cos n\theta = \frac{\sin(n\theta/2)}{\sin(\theta/2)} \cos\left(\frac{n+1}{2}\theta\right).$$

$$(ii) \sin \theta + \sin 2\theta + \sin 3\theta + \dots + \sin n\theta = \frac{\sin(n\theta/2)}{\sin(\theta/2)} \sin\left(\frac{n+1}{2}\theta\right).$$

Note : If $\theta = (2\pi/n)$ then the sum of the above series vanishes.

STRAIGHT LINES & CIRCLES IN TERMS OF COMPLEX NUMBERS :

(A) If z_1 & z_2 are two complex numbers then the complex number $z = \frac{nz_1 + mz_2}{m+n}$ divides the joins of z_1 & z_2 in the ratio $m:n$.

Note: (i) If a, b, c are three real numbers such that $az_1 + bz_2 + cz_3 = 0$;
where $a+b+c=0$ and a,b,c are not all simultaneously zero, then the complex numbers z_1, z_2 & z_3 are collinear.

(ii) If the vertices A, B, C of a Δ represent the complex nos. z_1, z_2, z_3 respectively, then :

$$(a) \text{Centroid of the } \Delta ABC = \frac{z_1 + z_2 + z_3}{3} :$$

$$(b) \text{Orthocentre of the } \Delta ABC = \frac{(a \sec A)z_1 + (b \sec B)z_2 + (c \sec C)z_3}{a \sec A + b \sec B + c \sec C} \text{ OR } \frac{z_1 \tan A + z_2 \tan B + z_3 \tan C}{\tan A + \tan B + \tan C}$$

$$(c) \text{Incentre of the } \Delta ABC = (az_1 + bz_2 + cz_3) \div (a + b + c).$$

$$(d) \text{Circumcentre of the } \Delta ABC = : (Z_1 \sin 2A + Z_2 \sin 2B + Z_3 \sin 2C) \div (\sin 2A + \sin 2B + \sin 2C).$$

amp(z) = θ is a ray emanating from the origin inclined at an angle θ to the x-axis.

$|z - a| = |z - b|$ is the perpendicular bisector of the line joining a to b .

The equation of a line joining z_1 & z_2 is given by ;

$$z = z_1 + t(z_1 - z_2) \text{ where } t \text{ is a parameter.}$$

$$z = z_1(1 + it) \text{ where } t \text{ is a real parameter is a line through the point } z_1 \text{ & perpendicular to } oz_1.$$

The equation of a line passing through z_1 & z_2 can be expressed in the determinant form as

$$\begin{vmatrix} z & \bar{z} & 1 \\ z_1 & \bar{z}_1 & 1 \\ z_2 & \bar{z}_2 & 1 \end{vmatrix} = 0. \text{ This is also the condition for three complex numbers to be collinear.}$$

Complex equation of a straight line through two given points z_1 & z_2 can be written as

$$z(\bar{z}_1 - \bar{z}_2) - \bar{z}(z_1 - z_2) + (z_1 \bar{z}_2 - \bar{z}_1 z_2) = 0, \text{ which on manipulating takes the form as } \bar{\alpha}z + \alpha\bar{z} + r = 0 \text{ where } r \text{ is real and } \alpha \text{ is a non zero complex constant.}$$

The equation of circle having centre z_0 & radius r is :

$$|z - z_0| = r \text{ or } z\bar{z} - z_0\bar{z} - \bar{z}_0z + \bar{z}_0z_0 - r^2 = 0 \text{ which is of the form}$$

$$z\bar{z} + \bar{\alpha}z + \alpha\bar{z} + r = 0, \text{ r is real centre } -\alpha \text{ & radius } \sqrt{\alpha\bar{\alpha} - r^2}.$$

Circle will be real if $\alpha\bar{\alpha} - r^2 \geq 0$.

The equation of the circle described on the line segment joining z_1 & z_2 as diameter is :

$$(i) \arg \frac{z - z_2}{z - z_1} = \pm \frac{\pi}{2} \text{ or } (z - z_1)(\bar{z} - \bar{z}_2) + (z - z_2)(\bar{z} - \bar{z}_1) = 0$$

(J) Condition for four given points z_1, z_2, z_3 & z_4 to be concyclic is, the number

$\frac{z_3 - z_1}{z_3 - z_2} \cdot \frac{z_4 - z_2}{z_4 - z_1}$ is real. Hence the equation of a circle through 3 non collinear points z_1, z_2 & z_3 can be taken as $\frac{(z-z_2)(z_3-z_1)}{(z-z_1)(z_3-z_2)}$ is real $\Rightarrow \frac{(z-z_2)(z_3-z_1)}{(z-z_1)(z_3-z_2)} = \frac{(\bar{z}-\bar{z}_2)(\bar{z}_3-\bar{z}_1)}{(\bar{z}-\bar{z}_1)(\bar{z}_3-\bar{z}_2)}$

13.(a) **Reflection points for a straight line :**

Two given points P & Q are the reflection points for a given straight line if the given line is the right bisector of the segment PQ. Note that the two points denoted by the complex numbers z_1 & z_2 will be the reflection points for the straight line $\bar{\alpha}z + \alpha\bar{z} + r = 0$ if and only if; $\bar{\alpha}z_1 + \alpha\bar{z}_2 + r = 0$, where r is real and α is non zero complex constant.

(b) **Inverse points w.r.t. a circle :**

Two points P & Q are said to be inverse w.r.t. a circle with centre 'O' and radius ρ , if :

(i) the point O, P, Q are collinear and on the same side of O. (ii) $OP \cdot OQ = \rho^2$.

Note that the two points z_1 & z_2 will be the inverse points w.r.t. the circle

$$z\bar{z} + \bar{\alpha}z + \alpha\bar{z} + r = 0 \text{ if and only if } z_1\bar{z}_2 + \bar{\alpha}z_1 + \alpha\bar{z}_2 + r = 0.$$

14. **PTOLEMY'S THEOREM :** It states that the product of the lengths of the diagonals of a convex quadrilateral inscribed in a circle is equal to the sum of the lengths of the two pairs of its opposite sides. i.e. $|z_1 - z_3| |z_2 - z_4| = |z_1 - z_2| |z_3 - z_4| + |z_1 - z_4| |z_2 - z_3|$.

15. **LOGARITHM OF A COMPLEX QUANTITY :**

(i) $\log_e(\alpha + i\beta) = \frac{1}{2} \log_e(\alpha^2 + \beta^2) + i \left(2n\pi + \tan^{-1} \frac{\beta}{\alpha} \right)$ where $n \in I$.

(ii) i^i represents a set of positive real numbers given by $e^{-\left(\frac{2n\pi+\pi}{2}\right)}$, $n \in I$.

VERY ELEMENTARY EXERCISE

Q.1 Simplify and express the result in the form of $a + bi$

(a) $\left(\frac{1+2i}{2+i}\right)^2$ (b) $-i(9+6i)(2-i)^{-1}$ (c) $\left(\frac{4i^3-i}{2i+1}\right)^2$ (d) $\frac{3+2i}{2-5i} + \frac{3-2i}{2+5i}$ (e) $\frac{(2+i)^2}{2-i} - \frac{(2-i)^2}{2+i}$

Q.2 Given that $x, y \in R$, solve : (a) $(x+2y)+i(2x-3y)=5-4i$ (b) $(x+iy)+(7-5i)=9+4i$

(c) $x^2 - y^2 - i(2x+y) = 2i$ (d) $(2+3i)x^2 - (3-2i)y = 2x - 3y + 5i$

(e) $4x^2 + 3xy + (2xy - 3x^2)i = 4y^2 - (x^2/2) + (3xy - 2y^2)i$

Q.3 Find the square root of : (a) $9 + 40i$ (b) $-11 - 60i$ (c) $50i$

Q.4 (a) If $f(x) = x^4 + 9x^3 + 35x^2 - x + 4$, find $f(-5+4i)$
 (b) If $g(x) = x^4 - x^3 + x^2 + 3x - 5$, find $g(2+3i)$

Q.5 Among the complex numbers z satisfying the condition $|z + 3 - \sqrt{3}i| = \sqrt{3}$, find the number having the least positive argument.

Q.6 Solve the following equations over C and express the result in the form $a + ib$, $a, b \in R$.

(a) $ix^2 - 3x - 2i = 0$ (b) $2(1+i)x^2 - 4(2-i)x - 5 - 3i = 0$

Q.7 Locate the points representing the complex number z on the Argand plane:

(a) $|z+1-2i| = \sqrt{7}$; (b) $|z-1|^2 + |z+1|^2 = 4$; (c) $\left|\frac{z-3}{z+3}\right| = 3$; (d) $|z-3| = |z-6|$

Q.8 If a & b are real numbers between 0 & 1 such that the points $z_1 = a+i$, $z_2 = 1+bi$ & $z_3 = 0$ form an equilateral triangle, then find the values of 'a' and 'b'.

Q.9 For what real values of x & y are the numbers $-3 + ix^2y$ & $x^2 + y + 4i$ conjugate complex?

Q.10 Find the modulus, argument and the principal argument of the complex numbers.

(i) $6(\cos 310^\circ - i \sin 310^\circ)$ (ii) $-2(\cos 30^\circ + i \sin 30^\circ)$ (iii) $\frac{2+i}{4i+(1+i)^2}$

Q.11 If $(x+iy)^{1/3} = a+bi$; prove that $4(a^2 - b^2) = \frac{x}{a} + \frac{y}{b}$.

Q.12 (a) If $\frac{a+ib}{c+id} = p+qi$, prove that $p^2 + q^2 = \frac{a^2 + b^2}{c^2 + d^2}$.

(b) Let z_1, z_2, z_3 be the complex numbers such that

$$z_1 + z_2 + z_3 = z_1z_2 + z_2z_3 + z_3z_1 = 0. \text{ Prove that } |z_1| = |z_2| = |z_3|.$$

Q.13 Let z be a complex number such that $z \in c\setminus R$ and $\frac{1+z+z^2}{1-z+z^2} \in R$, then prove that $|z|=1$.

Q.14 Prove the identity, $|1-z_1\bar{z}_2|^2 - |z_1 - z_2|^2 = (1-|z_1|^2)(1-|z_2|^2)$

- Q.15 For any two complex numbers, prove that $|z_1 + z_2|^2 + |z_1 - z_2|^2 = 2 [|z_1|^2 + |z_2|^2]$. Also give the geometrical interpretation of this identity.
- Q.16 (a) Find all non-zero complex numbers Z satisfying $\bar{Z} = iZ^2$.
 (b) If the complex numbers z_1, z_2, \dots, z_n lie on the unit circle $|z| = 1$ then show that $|z_1 + z_2 + \dots + z_n| = |z_1^{-1} + z_2^{-1} + \dots + z_n^{-1}|$.
- Q.17 Find the Cartesian equation of the locus of 'z' in the complex plane satisfying, $|z - 4| + |z + 4| = 16$.
- Q.18 If ω is an imaginary cube root of unity then prove that :
 (a) $(1 + \omega - \omega^2)^3 - (1 - \omega + \omega^2)^3 = 0$ (b) $(1 - \omega + \omega^2)^5 + (1 + \omega - \omega^2)^5 = 32$
 (c) If ω is the cube root of unity, Find the value of, $(1 + 5\omega^2 + \omega^4)(1 + 5\omega^4 + \omega^2)(5\omega^3 + \omega + \omega^2)$.
- Q.19 If ω is a cube root of unity, prove that ; (i) $(1 + \omega - \omega^2)^3 - (1 - \omega + \omega^2)^3$
 (ii) $\frac{a + b\omega + c\omega^2}{c + a\omega + b\omega^2} = \omega^2$ (iii) $(1 - \omega)(1 - \omega^2)(1 - \omega^4)(1 - \omega^8) = 9$
- Q.20 If $x = a + b$; $y = a\omega + b\omega^2$; $z = a\omega^2 + b\omega$, show that
 (i) $xyz = a^3 + b^3$ (ii) $x^2 + y^2 + z^2 = 6ab$ (iii) $x^3 + y^3 + z^3 = 3(a^3 + b^3)$
- Q.21 If $(w \neq 1)$ is a cube root of unity then $\begin{vmatrix} 1 & 1+i+w^2 & w^2 \\ 1-i & -1 & w^2-1 \\ -i & -i+w-1 & -1 \end{vmatrix} =$
 (A) 0 (B) 1 (C) i (D) w
- Q.22 (a) $(1 + w)^7 = A + Bw$ where w is the imaginary cube root of a unity and $A, B \in \mathbb{R}$, find the ordered pair (A, B) .
 (b) The value of the expression ;
 1. $(2 - w)(2 - w^2) + 2. (3 - w)(3 - w^2) + \dots + (n - 1) \cdot (n - w)(n - w^2)$, where w is an imaginary cube root of unity is _____.
- Q.23 If $n \in \mathbb{N}$, prove that $(1 + i)^n + (1 - i)^n = 2^{\frac{n}{2}+1} \cdot \cos \frac{n\pi}{4}$.
- Q.24 Show that the sum $\sum_{k=1}^{2n} \left(\sin \frac{2\pi k}{2n+1} - i \cos \frac{2\pi k}{2n+1} \right)$ simplifies to a pure imaginary number.
- Q.25 If $x = \cos \theta + i \sin \theta$ & $1 + \sqrt{1 - a^2} = na$, prove that $1 + a \cos \theta = \frac{a}{2n} (1 + nx) \left(1 + \frac{n}{x} \right)$.
- Q.26 The number t is real and not an integral multiple of $\pi/2$. The complex number x_1 and x_2 are the roots of the equation, $\tan^2(t) \cdot x^2 + \tan(t) \cdot x + 1 = 0$
 Show that $(x_1)^n + (x_2)^n = 2 \left(\cos \frac{2n\pi}{3} \right) \cot^n(t)$.

EXERCISE - 1

- Q.1 Simplify and express the result in the form of $a + bi$:
 (a) $-i(9 + 6i)(2 - i)^{-1}$ (b) $\left(\frac{4i^3 - i}{2i + 1} \right)^2$ (c) $\frac{3 + 2i}{2 - 5i} + \frac{3 - 2i}{2 + 5i}$
 (d) $\frac{(2+i)^2}{2-i} - \frac{(2-i)^2}{2+i}$ (e) $\sqrt{i} + \sqrt{-i}$
- Q.2 Find the modulus, argument and the principal argument of the complex numbers.
 (i) $z = 1 + \cos\left(\frac{10\pi}{9}\right) + i \sin\left(\frac{10\pi}{9}\right)$ (ii) $(\tan 1 - i)^2$
 (iii) $z = \frac{\sqrt{5+12i} + \sqrt{5-12i}}{\sqrt{5+12i} - \sqrt{5-12i}}$ (iv) $\frac{i-1}{i\left(1-\cos\frac{2\pi}{5}\right) + \sin\frac{2\pi}{5}}$
- Q.3 Given that $x, y \in \mathbb{R}$, solve :
 (a) $(x + 2y) + i(2x - 3y) = 5 - 4i$ (b) $\frac{x}{1+2i} + \frac{y}{3+2i} = \frac{5+6i}{8i-1}$
 (c) $x^2 - y^2 - i(2x + y) = 2i$ (d) $\frac{1+2i}{2+3i} x^2 - \frac{3+2i}{3-2i} y = 2x - 3y + 5i$
 (e) $4x^2 + 3xy + (2xy - 3x^2)i = 4y^2 - (x^2/2) + (3xy - 2y^2)i$
- Q.4(a) Let Z is complex satisfying the equation, $z^2 - (3 + i)z + m + 2i = 0$, where $m \in \mathbb{R}$.

Get Solution of These Packages & Learn by Video Tutorials on www.MathsBySuhag.com

Suppose the equation has a real root, then find the value of m .

(b) a, b, c are real numbers in the polynomial, $P(Z) = 2Z^4 + aZ^3 + bZ^2 + cZ + 3$

If two roots of the equation $P(Z) = 0$ are 2 and i , then find the value of ' a '.

Q.5(a) Find the real values of x & y for which $z_1 = 9y^2 - 4 - 10ix$ and

$$z_2 = 8y^2 - 20i \text{ are conjugate complex of each other.}$$

(b) Find the value of $x^4 - x^3 + x^2 + 3x - 5$ if $x = 2 + 3i$

o . 6 . Solve the following for z : (a) $z^2 - (3 - 2i)z = (5i - 5)$ (b) $|z| + z = 2 + i$

Q.7(a) If $iZ^3 + Z^2 - Z + i = 0$, then show that $|Z| = 1$.

(b) Let z_1 and z_2 be two complex numbers such that $\left| \frac{z_1 - 2z_2}{2 - z_1 \bar{z}_2} \right| = 1$ and $|z_2| \neq 1$, find $|z_1|$.

(c) Let $z_1 = 10 + 6i$ & $z_2 = 4 + 6i$. If z is any complex number such that the argument of, $\frac{z - z_1}{z - z_2}$ is $\frac{\pi}{4}$, then

Q.8 prove that $|z - 7 - 9i| = 3\sqrt{2}$.
Show that the product,

$$\left[1 + \left(\frac{1+i}{2} \right) \right] \left[1 + \left(\frac{1+i}{2} \right)^2 \right] \left[1 + \left(\frac{1+i}{2} \right)^2 \right]^2 \dots \left[1 + \left(\frac{1+i}{2} \right)^{2^n} \right] \text{ is equal to } \left(1 - \frac{1}{2^{2^n}} \right) (1+i) \text{ where } n \geq 2.$$

Q.9 Let a & b be complex numbers (which may be real) and let,

$$Z = z^3 + (a + b + 3i)z^2 + (ab + 3ia + 2ib - 2)z + 2abi - 2a.$$

(i) Show that Z is divisible by, $z + b + i$. (ii) Find all complex numbers z for which $Z = 0$.

(iii) Find all purely imaginary numbers a & b when $z = 1 + i$ and Z is a real number.

Q.10 Interpret the following locii in $z \in C$.

(a) $1 < |z - 2i| < 3$

(b) $\operatorname{Re} \left(\frac{z+2i}{iz+2} \right) \leq 4 \quad (z \neq 2i)$

(c) $\operatorname{Arg}(z+i) - \operatorname{Arg}(z-i) = \pi/2$

(d) $\operatorname{Arg}(z-a) = \pi/3$ where $a = 3 + 4i$.

Q.11 Prove that the complex numbers z_1 and z_2 and the origin form an isosceles triangle with vertical angle $2\pi/3$ if $z_1^2 + z_2^2 + z_1 z_2 = 0$.

Q.12 P is a point on the Aragand diagram. On the circle with OP as diameter two points Q & R are taken such that $\angle POQ = \angle QOR = \theta$. If 'O' is the origin & P, Q & R are represented by the complex numbers Z_1, Z_2 & Z_3 respectively, show that: $Z_2^2 \cdot \cos 2\theta = Z_1 \cdot Z_3 \cos^2 \theta$.

Q.13 Let z_1, z_2, z_3 are three pair wise distinct complex numbers and t_1, t_2, t_3 are non-negative real numbers such that $t_1 + t_2 + t_3 = 1$. Prove that the complex number $z = t_1 z_1 + t_2 z_2 + t_3 z_3$ lies inside a triangle with vertices z_1, z_2, z_3 or on its boundary.

Q.14 If a CiS α , b CiS β , c CiS γ represent three distinct collinear points in an Argand's plane, then prove the following :

(i) $\sum ab \sin(\alpha - \beta) = 0$.

(ii) $(a \operatorname{CiS} \alpha) \sqrt{b^2 + c^2 - 2bc \cos(\beta - \gamma)} \pm (b \operatorname{CiS} \beta) \sqrt{a^2 + c^2 - 2ac \cos(\alpha - \gamma)}$

$$\mp (c \operatorname{CiS} \gamma) \sqrt{a^2 + b^2 - 2ab \cos(\alpha - \beta)} = 0.$$

Q.15 Find all real values of the parameter a for which the equation $(a-1)z^4 - 4z^2 + a + 2 = 0$ has only pure imaginary roots.

Q.16 Let $A \equiv z_1; B \equiv z_2; C \equiv z_3$ are three complex numbers denoting the vertices of an acute angled triangle. If the origin 'O' is the orthocentre of the triangle, then prove that

$$z_1 \bar{z}_2 + \bar{z}_1 z_2 = z_2 \bar{z}_3 + \bar{z}_2 z_3 = z_3 \bar{z}_1 + \bar{z}_3 z_1$$

hence show that the ΔABC is a right angled triangle $\Leftrightarrow z_1 \bar{z}_2 + \bar{z}_1 z_2 = z_2 \bar{z}_3 + \bar{z}_2 z_3 = z_3 \bar{z}_1 + \bar{z}_3 z_1 = 0$

Q.17 If the complex number $P(w)$ lies on the standard unit circle in an Argand's plane and $z = (aw+b)(w-c)^{-1}$ then, find the locus of z and interpret it. Given a, b, c are real.

Q.18(a) Without expanding the determinant at any stage , find $K \in R$ such that

$$\begin{vmatrix} 4i & 8+i & 4+3i \\ -8+i & 16i & i \\ -4+Ki & i & 8i \end{vmatrix} \text{ has purely imaginary value.}$$

(b) If A, B and C are the angles of a triangle

$$D = \begin{vmatrix} e^{-2iA} & e^{iC} & e^{iB} \\ e^{iC} & e^{-2iB} & e^{iA} \\ e^{iB} & e^{iA} & e^{-2iC} \end{vmatrix} \text{ where } i = \sqrt{-1} \text{ then find the value of D.}$$

- Q.19 If w is an imaginary cube root of unity then prove that :
 (a) $(1 - w + w^2)(1 - w^2 + w^4)(1 - w^4 + w^8) \dots$ to $2n$ factors $= 2^{2n}$.
 (b) If w is a complex cube root of unity, find the value of
 $(1 + w)(1 + w^2)(1 + w^4)(1 + w^8) \dots$ to n factors .
- Q.20 Prove that $\left(\frac{1 + \sin \theta + i \cos \theta}{1 + \sin \theta - i \cos \theta} \right)^n = \cos\left(\frac{n\pi}{2} - n\theta\right) + i \sin\left(\frac{n\pi}{2} - n\theta\right)$. Hence deduce that
 $\left(1 + \sin \frac{\pi}{5} + i \cos \frac{\pi}{5}\right)^5 + i \left(1 + \sin \frac{\pi}{5} - i \cos \frac{\pi}{5}\right)^5 = 0$
- Q.21 If $\cos(\alpha - \beta) + \cos(\beta - \gamma) + \cos(\gamma - \alpha) = -3/2$ then prove that :
 (a) $\sum \cos 2\alpha = 0 = \sum \sin 2\alpha$ (b) $\sum \sin(\alpha + \beta) = 0 = \sum \cos(\alpha + \beta)$ (c) $\sum \sin^2 \alpha = \sum \cos^2 \alpha = 3/2$
 (d) $\sum \sin 3\alpha = 3 \sin(\alpha + \beta + \gamma)$ (e) $\sum \cos 3\alpha = 3 \cos(\alpha + \beta + \gamma)$
 (f) $\cos^3(\theta + \alpha) + \cos^3(\theta + \beta) + \cos^3(\theta + \gamma) = 3 \cos(\theta + \alpha) \cdot \cos(\theta + \beta) \cdot \cos(\theta + \gamma)$ where $\theta \in \mathbb{R}$.
- Q.22 Resolve $Z^5 + 1$ into linear & quadratic factors with real coefficients. Deduce that : $4 \cdot \sin \frac{\pi}{10} \cdot \cos \frac{\pi}{5} = 1$.
- Q.23 If $x = 1 + i\sqrt{3}$; $y = 1 - i\sqrt{3}$ & $z = 2$, then prove that $x^p + y^p = z^p$ for every prime $p > 3$.
- Q.24 If the expression $z^5 - 32$ can be factorised into linear and quadratic factors over real coefficients as
 $(z^5 - 32) = (z - 2)(z^2 - pz + 4)(z^2 - qz + 4)$ then find the value of $(p^2 + 2p)$.
- Q.25(a) Let $z = x + iy$ be a complex number, where x and y are real numbers. Let A and B be the sets defined by
 $A = \{z \mid |z| \leq 2\}$ and $B = \{z \mid (1 - i)z + (1 + i)\bar{z} \geq 4\}$. Find the area of the region $A \cap B$.
 (b) For all real numbers x , let the mapping $f(x) = \frac{1}{x-i}$, where $i = \sqrt{-1}$. If there exist real numbers a, b, c and d for which $f(a), f(b), f(c)$ and $f(d)$ form a square on the complex plane. Find the area of the square.
- ## EXERCISE - 2
- Q.1 If $\begin{vmatrix} p & q & r \\ q & r & p \\ r & p & q \end{vmatrix} = 0$; where p, q, r are the moduli of non-zero complex numbers u, v, w respectively,
 prove that, $\arg \frac{w}{v} = \arg \left(\frac{w-u}{v-u} \right)^2$.
- Q.2 The equation $x^3 = 9 + 46i$ where $i = \sqrt{-1}$ has a solution of the form $a + bi$ where a and b are integers.
 Find the value of $(a^3 + b^3)$.
- Q.3 Show that the locus formed by z in the equation $z^3 + iz = 1$ never crosses the co-ordinate axes in the Argand's plane. Further show that $|z| = \sqrt{\frac{-\operatorname{Im}(z)}{2\operatorname{Re}(z)\operatorname{Im}(z) + 1}}$
- Q.4 If ω is the fifth root of 2 and $x = \omega + \omega^2$, prove that $x^5 = 10x^2 + 10x + 6$.
- Q.5 Prove that, with regard to the quadratic equation $z^2 + (p + ip')z + q + iq' = 0$ where p, p', q, q' are all real.
 (i) if the equation has one real root then $q'^2 - pp'q' + qp'^2 = 0$.
 (ii) if the equation has two equal roots then $p^2 - p'^2 = 4q$ & $pp' = 2q'$.
 State whether these equal roots are real or complex.
- Q.6 If the equation $(z+1)^7 + z^7 = 0$ has roots z_1, z_2, \dots, z_7 , find the value of
 (a) $\sum_{r=1}^7 \operatorname{Re}(Z_r)$ and (b) $\sum_{r=1}^7 \operatorname{Im}(Z_r)$
- Q.7 Find the roots of the equation $Z^n = (Z+1)^n$ and show that the points which represent them are collinear on the complex plane. Hence show that these roots are also the roots of the equation
 $\left(2 \sin \frac{m\pi}{n}\right)^2 \bar{Z}^2 + \left(2 \sin \frac{m\pi}{n}\right)^2 \bar{Z} + 1 = 0$.
- Q.8 Dividing $f(z)$ by $z - i$, we get the remainder i and dividing it by $z + i$, we get the remainder

1 + i. Find the remainder upon the division of $f(z)$ by $z^2 + 1$.

- Q.9 Let z_1 & z_2 be any two arbitrary complex numbers then prove that :

$$|z_1 + z_2| \geq \frac{1}{2}(|z_1| + |z_2|) \left| \frac{z_1}{|z_1|} + \frac{z_2}{|z_2|} \right|.$$

- Q.10 If Z_r , $r = 1, 2, 3, \dots, 2m$, $m \in \mathbb{N}$ are the roots of the equation

$$Z^{2m} + Z^{2m-1} + Z^{2m-2} + \dots + Z + 1 = 0 \text{ then prove that } \sum_{r=1}^{2m} \frac{1}{Z_r - 1} = -m$$

- Q.11 If $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$ ($n \in \mathbb{N}$), prove that :

$$(a) C_0 + C_4 + C_8 + \dots = \frac{1}{2} \left[2^{n-1} + 2^{n/2} \cos \frac{n\pi}{4} \right] \quad (b) C_1 + C_5 + C_9 + \dots = \frac{1}{2} \left[2^{n-1} + 2^{n/2} \sin \frac{n\pi}{4} \right]$$

$$(c) C_2 + C_6 + C_{10} + \dots = \frac{1}{2} \left[2^{n-1} - 2^{n/2} \cos \frac{n\pi}{4} \right] \quad (d) C_3 + C_7 + C_{11} + \dots = \frac{1}{2} \left[2^{n-1} - 2^{n/2} \sin \frac{n\pi}{4} \right]$$

$$(e) C_0 + C_3 + C_6 + C_9 + \dots = \frac{1}{3} \left[2^n + 2 \cos \frac{n\pi}{3} \right]$$

- Q.12 Let z_1, z_2, z_3, z_4 be the vertices A, B, C, D respectively of a square on the Argand diagram taken in anticlockwise direction then prove that :

$$(i) 2z_2 = (1+i)z_1 + (1-i)z_3 \quad \& \quad (ii) 2z_4 = (1-i)z_1 + (1+i)z_3$$

- Q.13 Show that all the roots of the equation $\left(\frac{1+ix}{1-ix} \right)^n = \frac{1+ia}{1-ia}$ $a \in \mathbb{R}$ are real and distinct.

- Q.14 Prove that:

$$(a) \cos x + {}^nC_1 \cos 2x + {}^nC_2 \cos 3x + \dots + {}^nC_n \cos (n+1)x = 2^n \cdot \cos^n \frac{x}{2} \cdot \cos \left(\frac{n+2}{2}x \right)$$

$$(b) \sin x + {}^nC_1 \sin 2x + {}^nC_2 \sin 3x + \dots + {}^nC_n \sin (n+1)x = 2^n \cdot \cos^n \frac{x}{2} \cdot \sin \left(\frac{n+2}{2}x \right)$$

$$(c) \cos \left(\frac{2\pi}{2n+1} \right) + \cos \left(\frac{4\pi}{2n+1} \right) + \cos \left(\frac{6\pi}{2n+1} \right) + \dots + \cos \left(\frac{2n\pi}{2n+1} \right) = -\frac{1}{2} \text{ When } n \in \mathbb{N}.$$

- Q.15 Show that all roots of the equation $a_0z^n + a_1z^{n-1} + \dots + a_{n-1}z + a_n = n$,

where $|a_i| \leq 1$, $i = 0, 1, 2, \dots, n$ lie outside the circle with centre at the origin and radius $\frac{n-1}{n}$.

- Q.16 The points A, B, C depict the complex numbers z_1, z_2, z_3 respectively on a complex plane & the angle

B & C of the triangle ABC are each equal to $\frac{1}{2}(\pi - \alpha)$. Show that

$$(z_2 - z_3)^2 = 4(z_3 - z_1)(z_1 - z_2) \sin^2 \frac{\alpha}{2}.$$

- Q.17 Show that the equation $\frac{A_1^2}{x - a_1} + \frac{A_2^2}{x - a_2} + \dots + \frac{A_n^2}{x - a_n} = k$ has no imaginary root, given that:

$a_1, a_2, a_3, \dots, a_n$ & $A_1, A_2, A_3, \dots, A_n, k$ are all real numbers.

- Q.18 Let a, b, c be distinct complex numbers such that $\frac{a}{1-b} = \frac{b}{1-c} = \frac{c}{1-a} = k$. Find the value of k.

- Q.19 Let α, β be fixed complex numbers and z is a variable complex number such that,

$$|z - \alpha|^2 + |z - \beta|^2 = k.$$

Find out the limits for 'k' such that the locus of z is a circle. Find also the centre and radius of the circle.

- Q.20 C is the complex number. $f : \mathbb{C} \rightarrow \mathbb{R}$ is defined by $f(z) = |z^3 - z + 2|$. What is the maximum value of f on the unit circle $|z| = 1$?

- Q.21 Let $f(x) = \log_{\cos 3x} (\cos 2ix)$ if $x \neq 0$ and $f(0) = K$ (where $i = \sqrt{-1}$) is continuous at $x = 0$ then find the value of K. Use of L'Hospital's rule or series expansion not allowed.

- Q.22 If z_1, z_2 are the roots of the equation $az^2 + bz + c = 0$, with $a, b, c > 0$; $2b^2 > 4ac > b^2$; $z_1 \in$ third quadrant; $z_2 \in$ second quadrant in the argand's plane then, show that

$$\arg\left(\frac{z_1}{z_2}\right) = 2\cos^{-1}\left(\frac{b^2}{4ac}\right)^{1/2}$$

- Q.23 Find the set of points on the argand plane for which the real part of the complex number $(1+i)z^2$ is positive where $z = x + iy$, $x, y \in \mathbb{R}$ and $i = \sqrt{-1}$.
- Q.24 If a and b are positive integer such that $N = (a+ib)^3 - 107i$ is a positive integer. Find N .
- Q.25 If the biquadratic $x^4 + ax^3 + bx^2 + cx + d = 0$ ($a, b, c, d \in \mathbb{R}$) has 4 non real roots, two with sum $3+4i$ and the other two with product $13+i$. Find the value of ' b '.

EXERCISE - 3

- Q.1 Evaluate: $\sum_{p=1}^{32} (3p+2) \left(\sum_{q=1}^{10} \left(\sin \frac{2q\pi}{11} - i \cos \frac{2q\pi}{11} \right) \right)^p$. [REE '97, 6]
- Q.2(a) Let z_1 and z_2 be roots of the equation $z^2 + pz + q = 0$, where the co-efficients p and q may be complex numbers. Let A and B represent z_1 and z_2 in the complex plane. If $\angle AOB = \alpha \neq 0$ and $OA = OB$, where O is the origin. Prove that $p^2 = 4q \cos^2\left(\frac{\alpha}{2}\right)$. [JEE '97, 5]
- (b) Prove that $\sum_{k=1}^{n-1} (n-k) \cos \frac{2k\pi}{n} = -\frac{n}{2}$ where $n \geq 3$ is an integer. [JEE '97, 5]
- Q.3(a) If ω is an imaginary cube root of unity, then $(1+\omega-\omega^2)^7$ equals
 (A) 128ω (B) -128ω (C) $128\omega^2$
 (D) $-128\omega^2$ [JEE' 98, 2 + 2]
- (b) The value of the sum $\sum_{n=1}^{13} (i^n + i^{n+1})$, where $i = \sqrt{-1}$, equals
 (A) i (B) $i-1$ (C) $-i$
 (D) 0 [JEE' 98, 2 + 2]
- Q.4 Find all the roots of the equation $(3z-1)^4 + (z-2)^4 = 0$ in the simplified form of $a+ib$. [REE '98, 6]
- Q.5(a) If $i = \sqrt{-1}$, then $4 + 5 \left(-\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)^{334} + 3 \left(-\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)^{365}$ is equal to :
 (A) $1 - i\sqrt{3}$ (B) $-1 + i\sqrt{3}$ (C) $i\sqrt{3}$ (D) $-i\sqrt{3}$
- (b) For complex numbers z & ω , prove that, $|z|^2 \omega - |\omega|^2 z = z - \omega$ if and only if,
 $z = \omega$ or $z\bar{\omega} = 1$ [JEE '99, 2 + 10 (out of 200)]
- Q.6 If $\alpha = e^{\frac{2\pi i}{7}}$ and $f(x) = A_0 + \sum_{k=1}^{20} A_k x^k$, then find the value of,
 $f(x) + f(\alpha x) + \dots + f(\alpha^6 x)$ independent of α . [REE '99, 6]
- Q.7(a) If z_1, z_2, z_3 are complex numbers such that $|z_1| = |z_2| = |z_3| = \left[\frac{1}{z_1} + \frac{1}{z_2} + \frac{1}{z_3} \right] = 1$, then
 $|z_1 + z_2 + z_3|$ is :
 (A) equal to 1 (B) less than 1 (C) greater than 3 (D) equal to 3
- (b) If $\arg(z) < 0$, then $\arg(-z) - \arg(z) =$
 (A) π (B) $-\pi$ (C) $-\frac{\pi}{2}$ (D) $\frac{\pi}{2}$
 [JEE 2000 (Screening) 1 + 1 out of 35]
- Q.8 Given, $z = \cos \frac{2\pi}{2n+1} + i \sin \frac{2\pi}{2n+1}$, 'n' a positive integer, find the equation whose roots are,
 $\alpha = z + z^3 + \dots + z^{2n-1}$ & $\beta = z^2 + z^4 + \dots + z^{2n}$. [REE 2000 (Mains) 3 out of 100]
- Q.9(a) The complex numbers z_1, z_2 and z_3 satisfying $\frac{z_1 - z_3}{z_2 - z_3} = \frac{1 - i\sqrt{3}}{2}$ are the vertices of a triangle which is
 (A) of area zero (B) right-angled isosceles
 (C) equilateral (D) obtuse - angled isosceles

- (b) Let z_1 and z_2 be nth roots of unity which subtend a right angle at the origin. Then n must be of the form
 (A) $4k + 1$ (B) $4k + 2$ (C) $4k + 3$ (D) $4k$

[JEE 2001 (Scr) 1 + 1 out of 35]

- Q.10 Find all those roots of the equation $z^{12} - 56z^6 - 512 = 0$ whose imaginary part is positive.
 [REE 2000, 3 out of 100]

- Q.11(a) Let $\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$. Then the value of the determinant $\begin{vmatrix} 1 & 1 & 1 \\ 1 & -1-\omega^2 & \omega^2 \\ 1 & \omega^2 & \omega^4 \end{vmatrix}$ is

(A) 3ω (B) $3\omega(\omega-1)$ (C) $3\omega^2$ (D) $3\omega(1-\omega)$

- (b) For all complex numbers z_1, z_2 satisfying $|z_1| = 12$ and $|z_2 - 3 - 4i| = 5$, the minimum value of $|z_1 - z_2|$ is
 (A) 0 (B) 2 (C) 7 (D) 17

[JEE 2002 (Scr) 3+3]

- (c) Let a complex number $\alpha, \alpha \neq 1$, be a root of the equation

$z^{p+q} - z^p - z^q + 1 = 0$ where p, q are distinct primes.

Show that either $1 + \alpha + \alpha^2 + \dots + \alpha^{p-1} = 0$ or $1 + \alpha + \alpha^2 + \dots + \alpha^{q-1} = 0$, but not both together.
 [JEE 2002, (5)]

- Q.12(a) If z_1 and z_2 are two complex numbers such that $|z_1| < 1 < |z_2|$ then prove that $\left| \frac{1 - z_1 \bar{z}_2}{z_1 - z_2} \right| < 1$.

- (b) Prove that there exists no complex number z such that $|z| < \frac{1}{3}$ and $\sum_{r=1}^n a_r z^r = 1$ where $|a_r| < 2$.

[JEE-03, 2 + 2 out of 60]

- Q.13(a) ω is an imaginary cube root of unity. If $(1 + \omega^2)^m = (1 + \omega^4)^m$, then least positive integral value of m is
 (A) 6 (B) 5 (C) 4 (D) 3


[JEE 2004 (Scr)]

- (b) Find centre and radius of the circle determined by all complex numbers $z = x + iy$ satisfying $\left| \frac{(z - \alpha)}{(z - \beta)} \right| = k$,
 where $\alpha = \alpha_1 + i\alpha_2, \beta = \beta_1 + i\beta_2$ are fixed complex and $k \neq 1$.

[JEE 2004, 2 out of 60]
 $p(\sqrt{2}-1, \sqrt{2})$

- Q.14(a) The locus of z which lies in shaded region is best represented by

(A) $z : |z + 1| > 2, |\arg(z + 1)| < \pi/4$
 (B) $z : |z - 1| > 2, |\arg(z - 1)| < \pi/4$
 (C) $z : |z + 1| < 2, |\arg(z + 1)| < \pi/2$
 (D) $z : |z - 1| < 2, |\arg(z - 1)| < \pi/2$



- (b) If a, b, c are integers not all equal and w is a cube root of unity ($w \neq 1$), then the minimum value of $|a + bw + cw^2|$ is

(A) 0

(B) 1

(C) $\frac{\sqrt{3}}{2}$

(D) $\frac{1}{2}$

[JEE 2005 (Scr), 3 + 3]

- (c) If one of the vertices of the square circumscribing the circle $|z - 1| = \sqrt{2}$ is $2 + \sqrt{3}i$. Find the other vertices of square.

[JEE 2005 (Mains), 4]

- Q.15 If $w = \alpha + i\beta$ where $\beta \neq 0$ and $z \neq 1$, satisfies the condition that $\frac{w - \overline{w}z}{1 - z}$ is purely real, then the set of values of z is

(A) $\{z : |z| = 1\}$ (B) $\{z : z = \bar{z}\}$ (C) $\{z : z \neq 1\}$ (D) $\{z : |z| = 1, z \neq 1\}$

[JEE 2006, 3]

ANSWER KEY VERY ELEMENTARY EXERCISE

- Q.1 (a) $\frac{7}{25} + \frac{24}{25}i$; (b) $\frac{21}{5} - \frac{12}{5}i$; (c) $3 + 4i$; (d) $-\frac{8}{29} + 0i$; (e) $\frac{22}{5}i$

- Q.2 (a) $x = 1, y = 2$; (b) $(2, 9)$; (c) $(-2, 2)$ or $\left(-\frac{2}{3}, -\frac{2}{3}\right)$; (d) $(1, 1) \left(0, \frac{5}{2}\right)$ (e) $x = K, y = \frac{3K}{2}, K \in \mathbb{R}$

- Q.3** (a) $\pm(5+4i)$; (b) $\pm(5-6i)$ (c) $\pm 5(1+i)$
- Q.4** (a) -160 ; (b) $-(77+108i)$
- Q.5** $-\frac{3}{2} + \frac{3\sqrt{3}}{2}i$
- Q.6** (a) $-i, -2i$ (b) $\frac{3-5i}{2}$ or $-\frac{1+i}{2}$
- Q.7** (a) on a circle of radius $\sqrt{7}$ with centre $(-1, 2)$; (b) on a unit circle with centre at origin
(c) on a circle with centre $(-15/4, 0)$ & radius $9/4$; (d) a straight line
- Q.8** $a = b = 2 - \sqrt{3}$;
- Q.9** $x = 1, y = -4$ or $x = -1, y = -4$
- Q.10** (i) Modulus = 6, Arg = $2k\pi + \frac{5\pi}{18}$ ($K \in I$), Principal Arg = $\frac{5\pi}{18}$ ($K \in I$)
(ii) Modulus = 2, Arg = $2k\pi + \frac{7\pi}{6}$, Principal Arg = $-\frac{5\pi}{6}$
(iii) Modulus = $\frac{\sqrt{5}}{6}$, Arg = $2k\pi - \tan^{-1} 2$ ($K \in I$), Principal Arg = $-\tan^{-1} 2$
- Q.16** (a) $\frac{\sqrt{3}}{2} - \frac{i}{2}, -\frac{\sqrt{3}}{2} - \frac{i}{2}, i$; **Q.17** $\frac{x^2}{64} + \frac{y^2}{48} = 1$; **Q.18** (c) 64 ; **Q.21** A
- Q.22** (a) $(1, 1)$; (b) $\left[\frac{n(n+1)}{2}\right]^2 - n$

EXERCISE - 1

- Q.1** (a) $\frac{21}{5} - \frac{12}{5}i$ (b) $3+4i$ (c) $-\frac{8}{29} + 0i$ (d) $\frac{22}{5}i$ (e) $\pm\sqrt{2}+0i$ or $0\pm\sqrt{2}i$
- Q.2** (i) Principal Arg $z = -\frac{4\pi}{9}$; $|z| = 2 \cos \frac{4\pi}{9}$; Arg $z = 2k\pi - \frac{4\pi}{9}$ $k \in I$
(ii) Modulus = $\sec^2 1$, Arg = $2n\pi + (2 - \pi)$, Principal Arg = $(2 - \pi)$
(iii) Principal value of Arg $z = -\frac{\pi}{2}$ & $|z| = \frac{3}{2}$; Principal value of Arg $z = \frac{\pi}{2}$ & $|z| = \frac{2}{3}$
(iv) Modulus = $\frac{1}{\sqrt{2}} \csc \frac{\pi}{5}$, Arg $z = 2n\pi + \frac{11\pi}{20}$, Principal Arg = $\frac{11\pi}{20}$
- Q.3** (a) $x = 1, y = 2$; (b) $x = 1$ & $y = 2$; (c) $(-2, 2)$ or $\left(-\frac{2}{3}, -\frac{2}{3}\right)$; (d) $(1, 1) \left(0, \frac{5}{2}\right)$; (e) $x = K, y = \frac{3K}{2}$ $K \in R$
- Q.4** (a) 2, (b) $-11/2$ **Q.5** (a) $[-(2, 2); (-2, -2)]$ (b) $-(77+108i)$
- Q.6** (a) $z = (2+i)$ or $(1-3i)$; (b) $z = \frac{3+4i}{4}$
- Q.7** (b) 2
- Q.9** (ii) $z = -(b+i); -2i, -a$ (iii) $\left(-\frac{2ti}{3t+5}, ti\right)$ where $t \in R - \left\{-\frac{5}{3}\right\}$
- Q.10** (a) The region between the concentric circles with centre at $(0, 2)$ & radii 1 & 3 units
(b) region outside or on the circle with centre $\frac{1}{2} + 2i$ and radius $\frac{1}{2}$.
(c) semi circle (in the 1st & 4th quadrant) $x^2 + y^2 = 1$ (d) a ray emanating from the point $(3+4i)$ directed away from the origin & having equation $\sqrt{3}x - y + 4 - 3\sqrt{3} = 0$
- Q.15** $[-3, -2]$ **Q.17** $(1 - c^2)|z|^2 - 2(a + bc)(Re z) + a^2 - b^2 = 0$
- Q.18** (a) $K = 3$, (b) -4 **Q.19** (b) one if n is even; $-w^2$ if n is odd
- Q.22** $(Z+1)(Z^2 - 2Z \cos 36^\circ + 1)(Z^2 - 2Z \cos 108^\circ + 1)$ **Q.24** 4
- Q.25** (a) $\pi - 2$; (b) $1/2$

EXERCISE - 2

- Q.2** 35 **Q.6** (a) $-\frac{7}{2}$, (b) zero **Q.8** $\frac{iz}{2} + \frac{1}{2} + i$ **Q.18** $-\omega$ or $-\omega^2$
- Q.19** $k > \frac{1}{2} |\alpha - \beta|^2$ **Q.20** $|f(z)|$ is maximum when $z = \omega$, where ω is the cube root unity and $|f(z)| = \sqrt{13}$
- Q.21** $K = -\frac{4}{9}$

Q.23 required set is constituted by the angles without their boundaries, whose sides are the straight lines

$$y = (\sqrt{2} - 1)x \text{ and } y + (\sqrt{2} + 1)x = 0 \text{ containing the } x\text{-axis}$$

Q.24 198 **Q.25** 51

EXERCISE-3

Q.1 $48(1-i)$

Q.3 (a) D **(b)** B

$$\text{Q.4} \quad Z = \frac{(29+20\sqrt{2})+i(\pm 15+25\sqrt{2})}{82}, \quad \frac{(29-20\sqrt{2})+i(\pm 15-25\sqrt{2})}{82}$$

Q.5 (a) C

$$\text{Q.6} \quad 7A_0 + 7A_7x^7 + 7A_{14}x^{14} \quad \text{Q.7 (a) A (b) A} \quad \text{Q.8} \quad z^2 + z + \frac{\sin^2 n \theta}{\sin^2 \theta} = 0, \text{ where } \theta = \frac{2\pi}{2n+1}$$

Q.9 (a) C, **(b)** D

$$\text{Q.10} \quad \pm 1 + i\sqrt{3}, \frac{(\pm\sqrt{3}+i)}{\sqrt{2}}, \sqrt{2}i$$

Q.11 (a) B ; **(b)** B

$$\text{Q.13 (a) D ; (b) Centre} = \frac{k^2\beta - \alpha}{k^2 - 1}, \text{ Radius} = \frac{1}{(k^2 - 1)} \sqrt{|\alpha - k^2\beta|^2 - (k^2 \cdot |\beta|^2 - |\alpha|^2)(k^2 - 1)}$$

$$\text{Q.14 (a) A, (b) B, (c) } z_2 = -\sqrt{3}i; z_3 = (1-\sqrt{3})+i; z_4 = (1+\sqrt{3})-i$$

Q.15 D

EXERCISE-4

Part : (A) Only one correct option

1. If $|z| = 1$ and $\omega = \frac{z-1}{z+1}$ (where $z \neq -1$), the $\operatorname{Re}(\omega)$ is

(A) 0


(B) $-\frac{1}{|z+1|^2}$

(C) $\left| \frac{z}{z+1} \right| \cdot \frac{1}{|z+1|^2}$

(D) $\frac{\sqrt{2}}{|z+1|^2}$

[IIT – 2003, 3]

2. The locus of z which lies in shaded region (excluding the boundaries) is best represented by



[IIT – 2005, 3]

(A) $z : |z+1| > 2$ and $|\arg(z+1)| < \pi/4$ (B) $z : |z-1| > 2$ and $|\arg(z-1)| < \pi/4$
 (C) $z : |z+1| < 2$ and $|\arg(z+1)| < \pi/2$ (D) $z : |z-1| < 2$ and $|\arg(z+1)| < \pi/2$

3. If $w = \alpha + i\beta$, where $\beta \neq 0$ and $z \neq 1$, satisfies the condition that $\left(\frac{w - \bar{w}z}{1-z} \right)$ is purely real, then the set of values of z is

(A) $\{z : |z| = 1\}$

(B) $\{z : z = \bar{z}\}$

(C) $\{z : z \neq 1\}$

(D) $\{z : |z| = 1, z \neq 1\}$

[IIT – 2006, (3, -1)]

4. If $(\sqrt{3} + i)^{100} = 2^{99}(a + ib)$, then b is equal to

(A) $\sqrt{3}$

(B) $\sqrt{2}$

(C) 1

(D) none of these

5. If $\operatorname{Re}\left(\frac{z-8i}{z+6}\right) = 0$, then z lies on the curve

(A) $x^2 + y^2 + 6x - 8y = 0$

(B) $4x - 3y + 24 = 0$

(D) none of these

(C) $4ab$

6. If n_1, n_2 are positive integers then : $(1+i)^{n_1} + (1+i^3)^{n_1} + (1-i^5)^{n_2} + (1-i^7)^{n_2}$ is a real number if and only if

(A) $n_1 = n_2 + 1$

(B) $n_1 + 1 = n_2$

(C) $n_1 = n_2$

(D) n_1, n_2 are any two positive integers

7. The three vertices of a triangle are represented by the complex numbers, 0, z_1 and z_2 . If the triangle is equilateral, then

(A) $z_1^2 - z_2^2 = z_1 z_2$

(B) $z_2^2 - z_1^2 = z_1 z_2$

(C) $z_1^2 + z_2^2 = z_1 z_2$

(D) $z_1^2 + z_2^2 + z_1 z_2 = 0$

8. If $x^2 - x + 1 = 0$ then the value of $\sum_{n=1}^5 \left(x^n + \frac{1}{x^n} \right)^2$ is

(A) 8

(B) 10

(C) 12

(D) none of these

- FREE Download Study Package from website: www.TekoClasses.com & www.MathsBySuhag.com**
9. If α is nonreal and $\alpha = \sqrt[5]{1}$ then the value of $2^{|1+\alpha+\alpha^2+\alpha^{-2}-\alpha^{-1}|}$ is equal to
 (A) 4 (B) 2 (C) 1 (D) none of these
10. If $z = x + iy$ and $z^{1/3} = a - ib$ then $\frac{x}{a} - \frac{y}{b} = k(a^2 - b^2)$ where $k =$
 (A) 1 (B) 2 (C) 3 (D) 4
11. $\left[\frac{-1+i\sqrt{3}}{2}\right]^6 + \left[\frac{-1-i\sqrt{3}}{2}\right]^6 + \left[\frac{-1+i\sqrt{3}}{2}\right]^5 + \left[\frac{-1-i\sqrt{3}}{2}\right]^5$ is equal to :
 (A) 1 (B) -1 (C) 2 (D) none
12. Expressed in the form $r(\cos\theta + i\sin\theta)$, $-2 + 2i$ becomes :
 (A) $2\sqrt{2}\left[\cos\left(-\frac{\pi}{4}\right) + i\sin\left(-\frac{\pi}{4}\right)\right]$ (B) $2\sqrt{2}\left[\cos\left(\frac{3\pi}{4}\right) + i\sin\left(\frac{3\pi}{4}\right)\right]$
 (C) $2\sqrt{2}\left[\cos\left(-\frac{3\pi}{4}\right) + i\sin\left(-\frac{3\pi}{4}\right)\right]$ (D) $\sqrt{2}\left[\cos\left(-\frac{\pi}{4}\right) + i\sin\left(-\frac{\pi}{4}\right)\right]$
13. The number of solutions of the equation in z , $z\bar{z} - (3+i)z - (3-i)\bar{z} - 6 = 0$ is :
 (A) 0 (B) 1 (C) 2 (D) infinite
14. If $|z| = \max\{|z-1|, |z+1|\}$ then
 (A) $|z + \bar{z}| = \frac{1}{2}$ (B) $z + \bar{z} = 1$ (C) $|z + \bar{z}| = 1$ (D) none of these
15. If P, P' represent the complex number z_1 and its additive inverse respectively then the complex equation of the circle with PP' as a diameter is
 (A) $\frac{z}{z_1} = \left(\frac{z_1}{z}\right)$ (B) $z\bar{z} + z_1\bar{z}_1 = 0$ (C) $z\bar{z}_1 + \bar{z}z_1 = 0$ (D) none of these
16. The points $z_1 = 3 + \sqrt{3}i$ and $z_2 = 2\sqrt{3} + 6i$ are given on a complex plane. The complex number lying on the bisector of the angle formed by the vectors z_1 and z_2 is :
 (A) $z = \frac{(3+2\sqrt{3})}{2} + \frac{\sqrt{3}+2}{2}i$ (B) $z = 5 + 5i$
 (C) $z = -1 - i$ (D) none
17. The expression $\left[\frac{1+i\tan\alpha}{1-i\tan\alpha}\right]^n - \frac{1+i\tan n\alpha}{1-i\tan n\alpha}$ when simplified reduces to :
 (A) zero (B) $2\sin n\alpha$ (C) $2\cos n\alpha$ (D) none
18. All roots of the equation, $(1+z)^6 + z^6 = 0$:
 (A) lie on a unit circle with centre at the origin (B) lie on a unit circle with centre at $(-1, 0)$
 (C) lie on the vertices of a regular polygon with centre at the origin (D) are collinear
19. Points z_1 & z_2 are adjacent vertices of a regular octagon. The vertex z_3 adjacent to z_2 ($z_3 \neq z_1$) is represented by :
 (A) $z_2 + \frac{1}{\sqrt{2}}(1 \pm i)(z_1 + z_2)$ (B) $z_2 + \frac{1}{\sqrt{2}}(1 \pm i)(z_1 - z_2)$
 (C) $z_2 + \frac{1}{\sqrt{2}}(1 \pm i)(z_2 - z_1)$ (D) none of these
20. If $z = x + iy$ then the equation of a straight line $Ax + By + C = 0$ where $A, B, C \in \mathbb{R}$, can be written on the complex plane in the form $\bar{a}z + a\bar{z} + 2C = 0$ where 'a' is equal to :
 (A) $\frac{(A+iB)}{2}$ (B) $\frac{A-iB}{2}$ (C) $A + iB$ (D) none
21. The points of intersection of the two curves $|z-3| = 2$ and $|z| = 2$ in an argand plane are:
 (A) $\frac{1}{2}(7 \pm i\sqrt{3})$ (B) $\frac{1}{2}(3 \pm i\sqrt{7})$ (C) $\frac{3}{2} \pm i\sqrt{\frac{7}{2}}$ (D) $\frac{7}{2} \pm i\sqrt{\frac{3}{2}}$
22. The equation of the radical axis of the two circles represented by the equations, $|z-2| = 3$ and $|z-2-3i| = 4$ on the complex plane is :
 (A) $3iz - 3i\bar{z} - 2 = 0$ (B) $3iz - 3i\bar{z} + 2 = 0$ (C) $iz - i\bar{z} + 1 = 0$ (D) $2iz - 2i\bar{z} + 3 = 0$
23. If $\prod_{p=1}^r e^{ip\theta} = 1$ where \prod denotes the continued product, then the most general value of θ is :
 (A) $\frac{2n\pi}{r(r-1)}$ (B) $\frac{2n\pi}{r(r+1)}$ (C) $\frac{4n\pi}{r(r-1)}$ (D) $\frac{4n\pi}{r(r+1)}$
24. The set of values of $a \in \mathbb{R}$ for which $x^2 + i(a-1)x + 5 = 0$ will have a pair of conjugate imaginary roots is
 (A) \mathbb{R} (B) $\{1\}$ (C) $|a| a^2 - 2a + 21 > 0$ (D) none of these

25. If $|z_1 - 1| < 1$, $|z_2 - 2| < 2$, $|z_3 - 3| < 3$ then $|z_1 + z_2 + z_3|$
 (A) is less than 6 (B) is more than 3
 (C) is less than 12 (D) lies between 6 and 12
26. If $z_1, z_2, z_3, \dots, z_n$ lie on the circle $|z| = 2$, then the value of
 $E = |z_1 + z_2 + \dots + z_n| - 4 \left| \frac{1}{z_1} + \frac{1}{z_2} + \dots + \frac{1}{z_n} \right|$ is
 (A) 0 (B) n (C) -n (D) none of these
- Part : (B) May have more than one options correct
27. If z_1 lies on $|z| = 1$ and z_2 lies on $|z| = 2$, then
 (A) $3 \leq |z_1 - 2z_2| \leq 5$ (B) $1 \leq |z_1 + z_2| \leq 3$
 (C) $|z_1 - 3z_2| \geq 5$ (D) $|z_1 - z_2| \geq 1$
28. If z_1, z_2, z_3, z_4 are root of the equation $a_0 z^4 + z_1 z^3 + z_2 z^2 + z_3 z + z_4 = 0$, where a_0, a_1, a_2, a_3 and a_4 are real, then
 (A) $\bar{z}_1, \bar{z}_2, \bar{z}_3, \bar{z}_4$ are also roots of the equation (B) z_1 is equal to at least one of $\bar{z}_1, \bar{z}_2, \bar{z}_3, \bar{z}_4$
 (C) $-\bar{z}_1, -\bar{z}_2, -\bar{z}_3, -\bar{z}_4$ are also roots of the equation (D) none of these
29. If $a^3 + b^3 + 6abc = 8c^3$ & ω is a cube root of unity then :
 (A) a, c, b are in A.P. (B) a, c, b are in H.P.
 (C) $a + b\omega - 2c\omega^2 = 0$ (D) $a + b\omega^2 - 2c\omega = 0$
30. The points z_1, z_2, z_3 on the complex plane are the vertices of an equilateral triangle if and only if :
 (A) $\sum (z_1 - z_2)(z_2 - z_3) = 0$ (B) $z_1^2 + z_2^2 + z_3^2 = 2(z_1 z_2 + z_2 z_3 + z_3 z_1)$
 (C) $z_1^2 + z_2^2 + z_3^2 = z_1 z_2 + z_2 z_3 + z_3 z_1$ (D) $2(z_1^2 + z_2^2 + z_3^2) = z_1 z_2 + z_2 z_3 + z_3 z_1$
31. If $|z_1 + z_2| = |z_1 - z_2|$ then
 (A) $|\arg z_1 - \arg z_2| = \frac{\pi}{2}$ (B) $|\arg z_1 - \arg z_2| = \pi$
 (C) $\frac{z_1}{z_2}$ is purely real (D) $\frac{z_1}{z_2}$ is purely imaginary

EXERCISE-5

1. Given that $x, y \in \mathbb{R}$, solve : $4x^2 + 3xy + (2xy - 3x^2)i = 4y^2 - (x^2/2) + (3xy - 2y^2)i$
2. If α & β are any two complex numbers, prove that :
 $|\alpha - \sqrt{\alpha^2 - \beta^2}| + |\alpha + \sqrt{\alpha^2 - \beta^2}| = |\alpha + \beta| + |\alpha - \beta|$.
3. If α, β are the numbers between 0 and 1, such that the points $z_1 = \alpha + i$, $z_2 = 1 + \beta i$ and $z_3 = 0$ form an equilateral triangle, then find α and β .
4. ABCD is a rhombus. Its diagonals AC and BD intersect at the point M and satisfy $BD = 2AC$. If the points D and M represent the complex numbers $1+i$ and $2-i$ respectively, then find the complex number corresponding to A.
5. Show that the sum of the p^{th} powers of n^{th} roots of unity :
 (a) is zero, when p is not a multiple of n . (b) is equal to n , when p is a multiple of n .
6. If $(1+x)^n = p_0 + p_1 x + p_2 x^2 + p_3 x^3 + \dots$, then prove that :
 (a) $p_0 - p_2 + p_4 - \dots = 2^{n/2} \cos \frac{n\pi}{4}$ (b) $p_1 - p_3 + p_5 - \dots = 2^{n/2} \sin \frac{n\pi}{4}$
7. Prove that, $\log_e \left(\frac{1}{1 - e^{i\theta}} \right) = \log_e \left(\frac{1}{2} \operatorname{cosec} \frac{\theta}{2} \right) + i \left(\frac{\pi}{2} - \frac{\theta}{2} \right)$
8. If $i^{i^1} \dots^\infty = A + iB$, principal values only being considered, prove that
 (a) $\tan \frac{1}{2} \pi A = \frac{B}{A}$ (b) $A^2 + B^2 = e^{-\pi B}$
9. Prove that the roots of the equation, $(x-1)^n = x^n$ are $\frac{1}{2} \left(1 + i \cot \frac{r\pi}{n} \right)$, where
 $r = 0, 1, 2, \dots, (n-1)$ & $n \in \mathbb{N}$.
10. If $\cos(\alpha - \beta) + \cos(\beta - \gamma) + \cos(\gamma - \alpha) = -3/2$ then prove that :
 (a) $\sum \cos 2\alpha = 0 = \sum \sin 2\alpha$ (b) $\sum \sin(\alpha + \beta) = 0 = \sum \cos(\alpha + \beta)$
 (c) $\sum \sin 3\alpha = 3 \sin(\alpha + \beta + \gamma)$ (d) $\sum \cos 3\alpha = 3 \cos(\alpha + \beta + \gamma)$
 (e) $\sum \sin^2 \alpha = \sum \cos^2 \alpha = 3/2$
 (f) $\cos^3(\theta + \alpha) + \cos^3(\theta + \beta) + \cos^3(\theta + \gamma) = 3 \cos(\theta + \alpha) \cdot \cos(\theta + \beta) \cdot \cos(\theta + \gamma)$
 where $\theta \in \mathbb{R}$.

- Get Solution of These Packages & Learn by Video Tutorials on www.MathsBySuhag.com
11. If α, β, γ are roots of $x^3 - 3x^2 + 3x + 7 = 0$ (and ω is imaginary cube root of unity), then find the value of $\frac{\alpha - 1}{\beta - 1} + \frac{\beta - 1}{\gamma - 1} + \frac{\gamma - 1}{\alpha - 1}$.
 12. Given that, $|z - 1| = 1$, where 'z' is a point on the argand plane. Show that $\frac{z - 2}{z} = i \tan(\arg z)$.
 13. P is a point on the Argand diagram. On the circle with OP as diameter two points Q & R are taken such that $\angle POQ = \angle QOR = \theta$. If 'O' is the origin & P, Q & R are represented by the complex numbers Z_1, Z_2 & Z_3 respectively, show that : $Z_2^2 \cos 2\theta = Z_1 \cdot Z_3 \cos^2 \theta$.
 14. Find an expression for $\tan 7\theta$ in terms of $\tan \theta$, using complex numbers. By considering $\tan 7\theta = 0$, show that $x = \tan^2(3\pi/7)$ satisfies the cubic equation $x^3 - 21x^2 + 35x - 7 = 0$.
 15. If $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$ ($n \in \mathbb{N}$), prove that : $C_2 + C_6 + C_{10} + \dots = \frac{1}{2} \left[2^{n-1} - 2^{n/2} \cos \frac{n\pi}{4} \right]$
 16. Prove that : $\cos\left(\frac{2\pi}{2n+1}\right) + \cos\left(\frac{4\pi}{2n+1}\right) + \cos\left(\frac{6\pi}{2n+1}\right) + \dots + \cos\left(\frac{2n\pi}{2n+1}\right) = -\frac{1}{2}$ When $n \in \mathbb{N}$.
 17. Show that all the roots of the equation $a_1z^3 + a_2z^2 + a_3z + a_4 = 3$, where $|a_i| \leq 1$, $i = 1, 2, 3, 4$ lie outside the circle with centre origin and radius $2/3$.
 18. Prove that $\sum_{k=1}^{n-1} (n-k) \cos \frac{2k\pi}{n} = -\frac{n}{2}$, where $n \geq 3$ is an integer
 19. Show that the equation $\frac{A_1^2}{x - a_1} + \frac{A_2^2}{x - a_2} + \dots + \frac{A_n^2}{x - a_n} = k$ has no imaginary root, given that : $a_1, a_2, a_3, \dots, a_n$ & $A_1, A_2, A_3, \dots, A_n, k$ are all real numbers.
 20. Let z_1, z_2, z_3 be three distinct complex numbers satisfying, $\frac{1}{2}z_1 - 1\frac{1}{2} = \frac{1}{2}z_2 - 1\frac{1}{2} = \frac{1}{2}z_3 - 1\frac{1}{2}$. Let A, B & C be the points represented in the Argand plane corresponding to z_1, z_2 and z_3 resp. Prove that $z_1 + z_2 + z_3 = 3$ if and only if D ABC is an equilateral triangle.
 21. Let α, β be fixed complex numbers and z is a variable complex number such that, $|z - \alpha|^2 + |z - \beta|^2 = k$. Find out the limits for 'k' such that the locus of z is a circle. Find also the centre and radius of the circle.
 22. If $1, \alpha_1, \alpha_2, \alpha_3, \dots, \alpha_{n-1}$ are the n, n^{th} roots of unity, then prove that $(1 - \alpha_1)(1 - \alpha_2)(1 - \alpha_3)\dots(1 - \alpha_{n-1}) = n$. Hence prove that $\sin \frac{\pi}{n}, \sin \frac{2\pi}{n}, \sin \frac{3\pi}{n}, \dots, \sin \frac{(n-1)\pi}{n} = \frac{n}{2^{n-1}}$.
 23. Find the real values of the parameter 'a' for which at least one complex number $z = x + iy$ satisfies both the equality $|z - ai| = a + 4$ and the inequality $|z - 2| < 1$.
 24. Prove that, with regard to the quadratic equation $z^2 + (p + ip')z + q + iq' = 0$; where p, p', q, q' are all real.
 - if the equation has one real root then $q'^2 - pp'q' + qp'^2 = 0$.
 - if the equation has two equal roots then $p^2 - p'^2 = 4q$ & $pp' = 2q'$.
 State whether these equal roots are real or complex.
 25. The points A, B, C depict the complex numbers z_1, z_2, z_3 respectively on a complex plane & the angle B & C of the triangle ABC are each equal to $\frac{1}{2}(\pi - \alpha)$. Show that $(z_2 - z_3)^2 = 4(z_3 - z_1)(z_1 - z_2) \sin^2 \frac{\alpha}{2}$.
 26. If z_1, z_2 & z_3 are the affixes of three points A, B & C respectively and satisfy the condition $|z_1 - z_2| = |z_1| + |z_2|$ and $|(2-i)z_1 + iz_3| = |z_1| + |(1-i)z_1 + iz_3|$ then prove that ΔABC is a right angled.
 27. If $1, \alpha_1, \alpha_2, \alpha_3, \alpha_4$ be the roots of $x^5 - 1 = 0$, then prove that $\frac{\omega - \alpha_1}{\omega^2 - \alpha_1} \cdot \frac{\omega - \alpha_2}{\omega^2 - \alpha_2} \cdot \frac{\omega - \alpha_3}{\omega^2 - \alpha_3} \cdot \frac{\omega - \alpha_4}{\omega^2 - \alpha_4} = \omega$.
 28. If one the vertices of the square circumscribing the circle $|z - 1| = \sqrt{2}$ is $2 + \sqrt{3}i$. Find the other vertices of the square. [IIT – 2005, 4]

EXERCISE-4

1. A 2. C 3. D 4. A
5. A 6. D 7. C 8. A
9. A 11. D 12. A 13. B
14. D 15. D 16. A 17. B
18. A 19. D 20. C 21. C
22. B 23. B 24. D 25. B
26. C 27. A 28. ABCD 29. AB
30. ACD 31. AC 10. AD

EXERCISE-5

1. $x = K, y = \frac{3K}{2} \quad K \in \mathbb{R}$ 3. $2 - \sqrt{3}, 2 + \sqrt{3}$
4. $3 - \frac{i}{2}$ or $1 - \frac{3}{2}i$ 11. $3\omega^2$
21. $k > \frac{1}{2} |\alpha - \beta|^2$ 23. $\left(-\frac{21}{10}, -\frac{5}{6}\right)$
28. $-i\sqrt{3}, 1 - \sqrt{3} + i, 1 + \sqrt{3} - i$



For 39 Years Que. from IIT-JEE(Advanced) &
15 Years Que. from AIEEE (JEE Main)
we distributed a book in class room