

BINOMIAL THEOREM

Some questions (Assertion–Reason type) are given below. Each question contains **Statement – 1 (Assertion)** and **Statement – 2 (Reason)**. Each question has 4 choices (A), (B), (C) and (D) out of which **ONLY ONE** is correct. So select the correct choice : **Choices are :**

- (A) **Statement – 1** is True, **Statement – 2** is True; **Statement – 2** is a correct explanation for **Statement – 1**.
 (B) **Statement – 1** is True, **Statement – 2** is True; **Statement – 2** is NOT a correct explanation for **Statement – 1**.
 (C) **Statement – 1** is True, **Statement – 2** is False.
 (D) **Statement – 1** is False, **Statement – 2** is True.

BINOMIAL THEOREM

373. **Statement-1:** The binomial theorem provides an expansion for the expression $(a + b)^n$, where $a, b, n \in \mathbb{R}$.
Statement-2: All coefficients in a binomial expansion may be obtained by Pascal's triangle.
374. **Statement-1:** If n is an odd prime then integral part of $(\sqrt{5} + 2)^n - 2^{n+1}$ ($[x]$ is divisible by 20 n).
Statement-2: If n is prime then ${}^nC_1, {}^nC_2, {}^nC_3, \dots, {}^nC_{n-1}$ must be divisible by n .
375. **Statement-1 :** 2^{60} when divided by 7 leaves the remainder 1.
Statement-2 : $(1 + x)^n = 1 + n_1x$, where $n, n_1 \in \mathbb{N}$.
376. **Statement-1 :** ${}^{21}C_0 + {}^{21}C_1 + \dots + {}^{21}C_{10} = 2^{20}$
Statement-2 : ${}^{2n+1}C_0 + {}^{2n+1}C_1 + \dots + {}^{2n+1}C_{2n+1} = 2^{2n+1}$ and ${}^nC_r = {}^nC_{n-r}$
377. Let n be a positive integers and k be a whole number, $k \leq 2n$.
Statement-1 : The maximum value of ${}^{2n}C_k$ is ${}^{2n}C_n$.
Statement-2 : $\frac{{}^{2n}C_{k+1}}{{}^{2n}C_k} > 1$, for $k = 0, 1, 2, \dots, n - 1$.
378. Let n be a positive integer. **Statement-1 :** $3^{2n+2} - 8n - 9$ is divisible by 64.
Statement-2 : $3^{2n+2} - 8n - 9 = (1+8)^{n+1} - 8n - 9$ and in the binomial expansion of $(1+8)^{n+1}$, sum of first two terms is $8n + 9$ and after that each term is a multiple of 8^2 .
379. **Statement-1 :** If n is an odd prime, then integral part of $(\sqrt{5} + 2)^n$ is divisible by 20n.
Statement-2 : If n is prime, then ${}^nC_1, {}^nC_2, {}^nC_3 \dots, {}^nC_{n-1}$ must be divisible by n .
380. **Statement-1 :** The coefficient of x^{203} in the expression $(x - 1)(x^2 - 2)(x^2 - 3) \dots (x^{20} - 20)$ must be 13.
Statement-2 : The coefficient of x^8 in the expression $(2 + x)^2 (3 + x)^3 (4 + x)^4$ is equal to 30.
381. **Statement-1 :** $C_0^2 + C_1^2 + C_2^2 + C_3^2 + \dots + C_n^2 = \frac{2n!}{(n!)^2}$ **Statement-2 :** ${}^nC_0 - {}^nC_1 + {}^nC_2 - \dots + (-1)^n {}^nC_n = 0$
382. **Statement-1 :** Some of coefficient $(x - 2y + 4z)^n$ is 3^n
Statement-2 : Some of coefficient of $(c_0x_0 + c_1x_1 + c_2x_2 + \dots + c_nx_n)^n$ is 2^n
383. **Statement-1:** The greatest coefficient in the expansion of $(a_1 + a_2 + a_3 + a_4)^{17}$ is $\frac{17!}{(3!)^3 4!}$
Statement-2: The number of distinct terms in $(1 + x + x^2 + x^3 + x^4 + x^5)^{100}$ is 501.

384. **Statement-1:** The co-efficient of x^5 in the expansion of $(1+x^2)^5(1+x)^4$ is 120
Statement-2: The sum of the coefficients in the expansion of $(1+2x-3y+5z)^3$ is 125.
385. **Statement-1:** The number of distinct terms in $(1+x+x^2+x^3+x^4)^{1000}$ is 4001
Statement-2: The number of distinct terms in the expansion $(a_1+a_2+\dots+a_m)^n$ is ${}^{n+m-1}C_{m-1}$
386. **Statement-1:** In the expansion of $(1+x)^{30}$, greatest binomial coefficient is ${}^{30}C_{15}$
Statement-2: In the expansion of $(1+x)^{30}$, the binomial coefficients of equidistant terms from end & beginning are equal.
387. **Statement-1:** Integral part of $(\sqrt{3}+1)^{2n+1}$ is even where $n \in \mathbb{I}$.
Statement-2: Integral part of any integral power of the expression of the form of $p + \sqrt{q}$ is even.
388. **Statement-1:** $\sum_{r=4}^{20} {}^r C_4 = {}^{21}C_4$ **Statement-2:** $1+x+x^2+x^3+\dots+x^{n-1} = \frac{1-x^n}{1-x}$ = sum of n terms of GP.
389. **Statement-1:** Last two digits of the number $(13)^{41}$ are 31.
Statement-2: When a number is divided by 1000, the remainder gives the last three digits.
390. **Statement-1:** ${}^n C_0 + {}^n C_1 + {}^n C_2 + \dots + {}^n C_n = 2^n$ where $n \in \mathbb{N}$.
Statement-2: The all possible selections of n distinct objects are 2^n .
391. **Statement-1:** The integral part of $(5+2\sqrt{6})^n$ is odd, where $n \in \mathbb{N}$.
Statement-2: $(x+a)^n - (x-a)^n = 2[{}^n C_0 x^n + {}^n C_2 x^{n-2} a^2 + {}^n C_4 x^{n-4} a^4 + \dots]$
392. **Statement-1:** If n is even then ${}^{2n} C_1 + {}^{2n} C_3 + {}^{2n} C_5 + \dots + {}^{2n} C_{n-1} = 2^{2n-1}$
Statement-2: ${}^{2n} C_1 + {}^{2n} C_3 + {}^{2n} C_5 + \dots + {}^{2n} C_{2n-1} = 2^{2n-1}$
393. **Statement-1:** Any positive integral power of $(\sqrt{2}-1)$ can be expressed as $\sqrt{N}-\sqrt{N-1}$ for some natural number $N > 1$.
Statement-2: Any positive integral power of $\sqrt{2}-1$ can be expressed as $A+B\sqrt{2}$ where A and B are integers.
394. **Statement-1:** The term independent of x in the expansion of $\left(x + \frac{1}{x} + 3\right)^m$ is $\frac{4m!}{(2m!)^2}$.
Statement-2: The Coefficient of x^b in the expansion of $(1+x)^n$ is ${}^n C_b$.
395. **Statement-1:** The coefficient of x^8 in the expansion of $(1+3x+3x^2+x^3)^{17}$ is ${}^{51}C_2$.
Statement-2: Coefficient of x^r in the expansion of $(1+x)^n$ is ${}^n C_r$.
396. **Statement-1:** If $(1+x)^n = c_0+c_1x+c_2x^2+\dots+c_nx^n$ then
 $c_0-2.c_1+3.c_2-\dots+(-1)^n(n+1)c_n=0$
Statement-2: Coefficients of equidistant terms in the expansion of $(x+a)^n$ where $n \in \mathbb{N}$ are equal.
397. **Statement-1:** $\sum_{k=1}^n k \binom{n}{k}^2 = n \cdot {}^{2n-1}C_{n-1}$
Statement-2: If 2^{2003} is divided by 15 then remainder is 8.
398. **Statement-1:** The co-efficient of $(1+x^2)^5(1+x)^4$ is 120.
Statement-2: The integral part of $(\sqrt{5}+2)^{10}$ is odd.

ANSWER

373. D 374. A 375. A 376. A 377. A 378. A 379. A 380. C 381. B 382. C 383. D
 384. D 385. B 386. B 387. C 388. D 389. D 390. A 391. B 392. D 393. A 394. D 395. D
 396. B 397. B 398. D 399. A

QUE. FROM COMPT. EXAMS.

1. The value of $(\sqrt{2}+1)^6 + (\sqrt{2}-1)^6$ will be [RPET 1997]
 (a) -198 (b) 198 (c) 99 (d) -99
2. If $(1+ax)^n = 1+8x+24x^2+\dots$, then the value of a and n is [IIT 1983; Pb. CET 1994, 99]
 (a) 2, 4 (b) 2, 3 (c) 3, 6 (d) 1, 2
3. The coefficient of x^5 in the expansion of $(1+x^2)^5(1+x)^4$ is [EAMCET 1996; UPSEAT 2001; Pb. CET 2002]

BINOMIAL THEOREM PART 2 OF 2

- (a) 30 (b) 60 (c) 40 (d) None of these
4. If $\frac{(1-3x)^{1/2} + (1-x)^{5/3}}{\sqrt{4-x}}$ is approximately equal to $a+bx$ for small values of x , then $(a,b) =$
 (a) $\left(1, \frac{35}{24}\right)$ (b) $\left(1, -\frac{35}{24}\right)$ (c) $\left(2, \frac{35}{12}\right)$ (d) $\left(2, -\frac{35}{12}\right)$
5. The value of x in the expression $[x + x^{\log_{10}(x)}]^5$, if the third term in the expansion is 10,00,000 [Roorkee 1992]
 (a) 10 (b) 11 (c) 12 (d) None of these
6. If the coefficient of the middle term in the expansion of $(1+x)^{2n+2}$ is p and the coefficients of middle terms in the expansion of $(1+x)^{2n+1}$ are q and r , then
 (a) $p+q=r$ (b) $p+r=q$ (c) $p=q+r$ (d) $p+q+r=0$
7. In the polynomial $(x-1)(x-2)(x-3)\dots(x-100)$, the coefficient of x^{99} is [AMU 2002]
 (a) 5050 (b) -5050 (c) 100 (d) 99
8. The coefficient of x^{100} in the expansion of $\sum_{j=0}^{200} (1+x)^j$ is [UPSEAT 2004]
 (a) $\binom{200}{100}$ (b) $\binom{201}{102}$ (c) $\binom{200}{101}$ (d) $\binom{201}{100}$
9. If the coefficient of x^7 in $\left(ax^2 + \frac{1}{bx}\right)^{11}$ is equal to the coefficient of x^{-7} in $\left(ax - \frac{1}{bx^2}\right)^{11}$, then $ab =$ [MP PET 1999; AMU 2001; Pb. CET 2002; AIEEE 2005]
 (a) 1 (b) 1/2 (c) 2 (d) 3
10. If the coefficient of x in the expansion of $\left(x^2 + \frac{k}{x}\right)^5$ is 270, then $k =$ [EAMCET 2002]
 (a) 1 (b) 2 (c) 3 (d) 4
11. The coefficients of three successive terms in the expansion of $(1+x)^n$ are 165, 330 and 462 respectively, then the value of n will be [UPSEAT 1999]
 (a) 11 (b) 10 (c) 12 (d) 8
12. If the coefficient of $(2r+4)^{th}$ and $(r-2)^{th}$ terms in the expansion of $(1+x)^{18}$ are equal, then $r =$ [MP PET 1997; Pb. CET 2001]
 (a) 12 (b) 10 (c) 8 (d) 6
13. The middle term in the expansion of $(1+x)^{2n}$ is [Pb. CET 1998]
 (a) $\frac{1.3.5\dots(5n-1)}{n!} x^n$ (b) $\frac{2.4.6\dots 2n}{n!} x^{2n+1}$ (c) $\frac{1.3.5\dots(2n-1)}{n!} x^n$ (d) $\frac{1.3.5\dots(2n-1)}{n!} 2^n x^n$
14. The value of $\binom{30}{0}\binom{30}{10} - \binom{30}{1}\binom{30}{11} + \binom{30}{2}\binom{30}{12} + \dots + \binom{30}{20}\binom{30}{30}$ [IIT Screening 2005]
 (a) ${}^{60}C_{20}$ (b) ${}^{30}C_{10}$ (c) ${}^{60}C_{30}$ (d) ${}^{40}C_{30}$
15. Middle term in the expansion of $(1+3x+3x^2+x^3)^6$ is [MP PET 1997]
 (a) 4^{th} (b) 3^{rd} (c) 10^{th} (d) None of these
16. Two middle terms in the expansion of $\left(x - \frac{1}{x}\right)^{11}$ are
 (a) $231x$ and $\frac{231}{x}$ (b) $462x$ and $\frac{462}{x}$ (c) $-462x$ and $\frac{462}{x}$ (d) None of these

BINOMIAL THEOREM PART 2 OF 2

17. The term independent of y in the expansion of $(y^{-1/6} - y^{1/3})^9$ is [BIT Ranchi 1980]
 (a) 84 (b) 8.4 (c) 0.84 (d) -84
18. The coefficient of the term independent of x in the expansion of $(1 + x + 2x^3)\left(\frac{3}{2}x^2 - \frac{1}{3x}\right)^9$ is [DCE 1994]
 (a) $\frac{1}{3}$ (b) $\frac{19}{54}$ (c) $\frac{17}{54}$ (d) $\frac{1}{4}$
19. The term independent of x in $\left[\frac{\sqrt{x}}{3} + \frac{\sqrt{3}}{x^2}\right]^{10}$ is [EAMCET 1984; RPET 2000]
 (a) $\frac{2}{3}$ (b) $\frac{5}{3}$ (c) $\frac{4}{3}$ (d) None of these
20. The term independent of x in $\left(\sqrt{x} - \frac{2}{x}\right)^{18}$ is [EAMCET 1990]
 (a) ${}^{18}C_6 2^6$ (b) ${}^{18}C_6 2^{12}$ (c) ${}^{18}C_{18} 2^{18}$ (d) None of these
21. The largest term in the expansion of $(3 + 2x)^{50}$ where $x = \frac{1}{5}$ is [IIT Screening 1993]
 (a) 5th (b) 51st (c) 7th (d) 6th
22. $\frac{C_1}{C_0} + 2\frac{C_2}{C_1} + 3\frac{C_3}{C_2} + \dots + 15\frac{C_{15}}{C_{14}} =$ [IIT 1962]
 (a) 100 (b) 120 (c) -120 (d) None of these
23. $\binom{n}{0} + 2\binom{n}{1} + 2^2\binom{n}{2} + \dots + 2^n\binom{n}{n}$ is equal to [AMU 2000]
 (a) 2^n (b) 0 (c) 3^n (d) None of these
24. If C_r stands for nC_r , the sum of the given series $\frac{2(n/2)!(n/2)!}{n!}[C_0^2 - 2C_1^2 + 3C_2^2 - \dots + (-1)^n(n+1)C_n^2]$, Where n is an even positive integer, is [IIT 1986]
 (a) 0 (b) $(-1)^{n/2}(n+1)$ (c) $(-1)^n(n+2)$ (d) $(-1)^{n/2}(n+2)$
25. Sum of odd terms is A and sum of even terms is B in the expansion $(x + a)^n$, then [RPET 1987; UPSEAT 2004]
 (a) $AB = \frac{1}{4}(x - a)^{2n} - (x + a)^{2n}$ (b) $2AB = (x + a)^{2n} - (x - a)^{2n}$
 (c) $4AB = (x + a)^{2n} - (x - a)^{2n}$ (d) None of these
26. In the expansion of $(x + a)^n$, the sum of odd terms is P and sum of even terms is Q , then the value of $(P^2 - Q^2)$ will be [RPET 1997; Pb. CET 1998]
 (a) $(x^2 + a^2)^n$ (b) $(x^2 - a^2)^n$ (c) $(x - a)^{2n}$ (d) $(x + a)^{2n}$
27. The sum of the coefficients in the expansion of $(1 + x - 3x^2)^{2163}$ will be [IIT 1982]
 (a) 0 (b) 1 (c) -1 (d) 2^{2163}
28. If the sum of the coefficients in the expansion of $(1 - 3x + 10x^2)^n$ is a and if the sum of the coefficients in the expansion of $(1 + x^2)^n$ is b , then [UPSEAT 2001]
 (a) $a = 3b$ (b) $a = b^3$ (c) $b = a^3$ (d) None of these
29. The sum of the coefficients in the expansion of $(x + y)^n$ is 4096. The greatest coefficient in the expansion is [Kurukshestra CEE 1998; AIEEE 2002]

BINOMIAL THEOREM PART 2 OF 2

- (a) 1024 (b) 924 (c) 824 (d) 724
30. If the sum of the coefficients in the expansion of $(\alpha x^2 - 2x + 1)^{35}$ is equal to the sum of the coefficients in the expansion of $(x - \alpha y)^{35}$, then $\alpha =$
- (a) 0 (b) 1
(c) May be any real number (d) No such value exist
31. For every natural number n , $3^{2n+2} - 8n - 9$ is divisible by
- (a) 16 (b) 128 (c) 256 (d) None of these [IIT 1977]
32. The least remainder when 17^{30} is divided by 5 is
- (a) 1 (b) 2 (c) 3 (d) 4 [Karnataka CET 2003]
33. The value of the natural numbers n such that the inequality $2^n > 2n + 1$ is valid is
- (a) For $n \geq 3$ (b) For $n < 3$ (c) For mn (d) For any n [MNR 1994]
34. Let $P(n)$ be a statement and let $P(n) \Rightarrow P(n + 1)$ for all natural numbers n , then $P(n)$ is true
- (a) For all n (b) For all $n > 1$
(c) For all $n > m$, m being a fixed positive integer
(d) Nothing can be said
35. $(1 + x)^n - nx - 1$ is divisible by (where $n \in N$)
- (a) $2x$ (b) x^2 (c) $2x^3$ (d) All of these

ANSWER KEY

1	b	2	a	3	b	4	b	5	a
6	c	7	b	8	a	9	a	10	c
11	a	12	d	13	d	14	b	15	c
16	c	17	d	18	c	19	b	20	a
21	c	22	b	23	c	24	d	25	c
26	b	27	c	28	b	29	b	30	b
31	a	32	d	33	a	34	d	35	b

**for 39 Yrs. Que. of IIT-JEE
&
15 Yrs. Que. of AIEEE
we have distributed already a book**