

Exercise - 7**Part : (A) Only one correct option**

1. If $a, b, c > 0$ & $x, y, z \in \mathbb{R}$ then the determinant $\begin{vmatrix} (a^x + a^{-x})^2 & (a^x - a^{-x})^2 & 1 \\ (b^y + b^{-y})^2 & (b^y - b^{-y})^2 & 1 \\ (c^z + c^{-z})^2 & (c^z - c^{-z})^2 & 1 \end{vmatrix} =$
- (A) $a^x b^y c^z$ (B) $a^{-x} b^{-y} c^{-z}$ (C) $a^{2x} b^{2y} c^{2z}$ (D) zero
2. If $a, b & c$ are non-zero real numbers then $D = \begin{vmatrix} b^2 c^2 & bc & b+c \\ c^2 a^2 & ca & c+a \\ a^2 b^2 & ab & a+b \end{vmatrix} =$
- (A) abc (B) $a^2 b^2 c^2$ (C) $bc + ca + ab$ (D) zero
3. The determinant $\begin{vmatrix} b_1 + c_1 & c_1 + a_1 & a_1 + b_1 \\ b_2 + c_2 & c_2 + a_2 & a_2 + b_2 \\ b_3 + c_3 & c_3 + a_3 & a_3 + b_3 \end{vmatrix} =$
- (A) $\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$ (B) $2 \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$ (C) $3 \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$ (D) none of these
4. The system of linear equations $x + y - z = 6$, $x + 2y - 3z = 14$ and $2x + 5y - \lambda z = 9$ ($\lambda \in \mathbb{R}$) has a unique solution if
- (A) $\lambda = 8$ (B) $\lambda \neq 8$ (C) $\lambda = 7$ (D) $\lambda \neq 7$
5. If the system of equations $x + 2y + 3z = 4$, $x + py + 2z = 3$, $x + 4y + \mu z = 3$ has an infinite number of solutions then:
- (A) $p = 2, \mu = 3$ (B) $p = 2, \mu = 4$ (C) $3p = 2\mu$ (D) none of these
6. Let $f(\theta) = \begin{vmatrix} \cos^2 \theta & \cos \theta \sin \theta & -\sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta & \cos \theta \\ \sin \theta & -\cos \theta & 0 \end{vmatrix}$ then $f\left(\frac{\pi}{6}\right) =$
- (A) 0 (B) 1 (C) 2 (D) none
7. The determinant $\begin{vmatrix} \cos(\theta+\phi) & -\sin(\theta+\phi) & \cos 2\phi \\ \sin \theta & \cos \theta & \sin \phi \\ -\cos \theta & \sin \theta & \cos \phi \end{vmatrix}$ is:
- (A) 0 (B) independent of θ (C) independent of ϕ (D) independent of θ & ϕ both
8. Value of $\Delta = \begin{vmatrix} \sin(2\alpha) & \sin(\alpha + \beta) & \sin(\alpha + \gamma) \\ \sin(\beta + \alpha) & \sin(2\beta) & \sin(\gamma + \beta) \\ \sin(\gamma + \alpha) & \sin(\gamma + \beta) & \sin(2\gamma) \end{vmatrix}$ is
- (A) $\Delta = 0$ (B) $\Delta = \sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma$
 (C) $\Delta = 3/2$ (D) none of these
9. If a, b, c are complex numbers and $z = \begin{vmatrix} 0 & -b & -c \\ b & 0 & -a \\ c & a & 0 \end{vmatrix}$ is
- (A) purely real (B) purely imaginary (C) 0 (D) none of these
10. If A, B, C are angles of a triangle ABC, then $\begin{vmatrix} \sin \frac{A}{2} & \sin \frac{B}{2} & \sin \frac{C}{2} \\ \sin(A+B+C) & \sin \frac{B}{2} & \cos \frac{A}{2} \\ \cos \frac{(A+B+C)}{2} & \tan(A+B+C) & \sin \frac{C}{2} \end{vmatrix}$ is less than or equal to
- (A) $\frac{3\sqrt{3}}{8}$ (B) $\frac{1}{8}$ (C) $2\sqrt{2}$ (D) 2

11. $\Delta = \begin{vmatrix} 1 & \frac{4\sin B}{b} & \cos A \\ 2a & 8\sin A & 1 \\ 3a & 12\sin A & \cos B \end{vmatrix}$ is (where a, b, c are the sides opposite to angles A, B, C respectively in a triangle)
- (A) $\frac{1}{2} \cos 2A$ (B) 0 (C) $\frac{1}{2} \sin 2A$ (D) $\frac{1}{2} (\cos^2 A + \cos^2 B)$
12. If $\begin{vmatrix} (b+c)^2 & a^2 & a^2 \\ b^2 & (c+a)^2 & b^2 \\ c^2 & c^2 & (a+b)^2 \end{vmatrix} = k abc (a+b+c)^3$ then the value of k is
- (A) 1 (B) 2 (C) 0 (D) $ab + bc + ac$
13. Let m be a positive integer & $D_r = \begin{vmatrix} 2r-1 & {}^m C_r & 1 \\ m^2-1 & 2^m & m+1 \\ \sin^2(m^2) & \sin^2(m) & \sin^2(m+1) \end{vmatrix}$ ($0 \leq r \leq m$) then the value of $\sum_{r=0}^m D_r$ is given by:
- (A) 0 (B) $m^2 - 1$ (C) 2^m (D) $2^m \sin^2(2^m)$
14. If a, b, c, are real numbers, and $D = \begin{vmatrix} a & 1+2i & 3-5i \\ 1-2i & b & -7-3i \\ 3+5i & -7+3i & c \end{vmatrix}$ then D is
- (A) purely real (B) purely imaginary (C) non real (D) integer
15. If $f(x) = \begin{vmatrix} 1 & x & x+1 \\ 2x & x(x-1) & (x+1)x \\ 3x(x-1) & x(x-1)(x-2) & (x+1)x(x-1) \end{vmatrix}$ then $f(100)$ is equal to:
- (A) 0 (B) 1 (C) 100 (D) -100
- Part : (B) May have more than one options correct**
16. Let $\phi_1(x) = x + a_1$, $\phi_2(x) = x^2 + b_1x + b_2$ and $\Delta = \begin{vmatrix} \phi_1(x_1) & \phi_1(x_2) & \phi_1(x_3) \\ \phi_2(x_1) & \phi_2(x_2) & \phi_2(x_3) \end{vmatrix}$, then
- (A) Δ is independent of a_1
 (C) Δ is independent of x_1, x_2 and x_3
- (B) Δ is independent of b_1 and b_2
 (D) none of these
17. If $\Delta = \begin{vmatrix} x & 2y-z & -z \\ y & 2x-z & -z \\ y & 2y-z & 2x-2y-z \end{vmatrix}$, then
- (A) $x-y$ is a factor of Δ
 (C) $(x-y)^3$ is a factor of Δ
- (B) $(x-y)^2$ is a factor of Δ
 (D) Δ is independent of z
18. Let $\Delta = \begin{vmatrix} \sin \theta \cos \phi & \sin \theta \sin \phi & \cos \theta \\ \cos \theta \cos \phi & \cos \theta \sin \phi & -\sin \theta \\ -\sin \theta \sin \phi & \sin \theta \cos \phi & 0 \end{vmatrix}$, then
- (A) Δ is independent of θ
 (C) Δ is a constant
- (B) Δ is independent of ϕ
 (D) $\left[\frac{d\Delta}{d\theta} \right]_{\theta=\pi/2} = 0$
19. Let $\Delta = \begin{vmatrix} a & -1 & 0 \\ ax & a & -1 \\ ax^2 & ax & a \end{vmatrix}$, then
- (A) $x+a$ is a factor of Δ
 (C) $(x+a)^3$ is a factor of Δ
- (B) $(x+a)^2$ is a factor of Δ
 (D) $(x+a)^4$ is not a factor of Δ
20. Let $\Delta = \begin{vmatrix} 1 & x & x^2 \\ x^2 & 1 & x \\ x & x^2 & 1 \end{vmatrix}$, then

Get Solution of These Packages & Learn by Video Tutorials on www.MathsBySuhag.com

- (A) $1 - x^3$ is a factor of Δ
 (C) $\Delta(x) = 0$ has 4 real roots

- (B) $(1 - x^3)^2$ is factor of Δ
 (D) $\Delta'(1) = 0$

21.

The determinant $\Delta = \begin{vmatrix} b & c & b\alpha + c \\ c & d & c\alpha + d \\ b\alpha + c & c\alpha + d & a\alpha^3 - ca \end{vmatrix}$ is equal to zero if

- (A) b, c, d are in A.P.
 (C) b, c, d are in H.P.
 (B) b, c, d are in G.P.
 (D) α is a root of $ax^3 - bx^2 - 3cx - d = 0$

Exercise - 8

1.

Using the properties of determinants, evaluate:

$$(i) \begin{vmatrix} 103 & 115 & 114 \\ 111 & 108 & 106 \\ 104 & 113 & 116 \end{vmatrix} + \begin{vmatrix} 113 & 116 & 104 \\ 108 & 106 & 111 \\ 115 & 114 & 103 \end{vmatrix}. \quad (ii) \begin{vmatrix} \sqrt{13} + \sqrt{3} & 2\sqrt{5} & \sqrt{5} \\ \sqrt{15} + \sqrt{26} & 5 & \sqrt{10} \\ 3 + \sqrt{65} & \sqrt{15} & 5 \end{vmatrix}.$$

2.

Find the non-zero roots of the equation, $\Delta = \begin{vmatrix} a & b & ax + b \\ b & c & bx + c \\ ax + b & bx + c & c \end{vmatrix} = 0$.

3.

Show that $\Delta = \begin{vmatrix} b^2 + c^2 & ab & ac \\ ab & c^2 + a^2 & bc \\ ca & cb & a^2 + b^2 \end{vmatrix} = 4a^2b^2c^2$

4.

Prove that, $\begin{vmatrix} 2 & \alpha + \beta + \gamma + \delta & \alpha\beta + \gamma\delta \\ \alpha + \beta + \gamma + \delta & 2(\alpha + \beta)(\gamma + \delta) & \alpha\beta(\gamma + \delta) + \gamma\delta(\alpha + \beta) \\ \alpha\beta + \gamma\delta & \alpha\beta(\gamma + \delta) + \gamma\delta(\alpha + \beta) & 2\alpha\beta\gamma\delta \end{vmatrix} = 0$.

5.

If $S_r = \alpha^r + \beta^r + \gamma^r$ then show that $\begin{vmatrix} S_0 & S_1 & S_2 \\ S_1 & S_2 & S_3 \\ S_2 & S_3 & S_4 \end{vmatrix} = (\alpha - \beta)^2 (\beta - \gamma)^2 (\gamma - \alpha)^2$.

6.

Find the value of 'a' if the three equations, $(a+1)^3x + (a+2)^3y = (a+3)^3$; $(a+1)x + (a+2)y = (a+3)$ & $x+y=1$ are consistent.

7.

Investigate for what values of λ, μ the simultaneous equations $x + y + z = 6$; $x + 2y + 3z = 10$ & $x + 2y + \lambda z = \mu$ have;

- (a) A unique solution
 (b) An infinite number of solutions.
 (c) No solution.

8.

Find those values of c for which the equations:

$$\begin{aligned} 2x + 3y &= 3 \\ (c+2)x + (c+4)y &= c+6 \\ (c+2)^2x + (c+4)^2y &= (c+6)^2 \end{aligned}$$

are consistent.

Also solve above equations for these values of c.

9.

Prove that $\Delta = \begin{vmatrix} \beta\gamma & \beta\gamma' + \beta'\gamma & \beta'\gamma' \\ \gamma\alpha & \gamma\alpha' + \gamma'\alpha & \gamma'\alpha' \\ \alpha\beta & \alpha\beta' + \alpha'\beta & \alpha'\beta' \end{vmatrix} = (\alpha\beta' - \alpha'\beta) (\beta\gamma' - \beta'\gamma) (\gamma\alpha' - \gamma'\alpha)$

10.

If $a^2 + b^2 + c^2 = 1$, then prove that $\begin{vmatrix} a^2 + (b^2 + c^2)\cos\phi & ab(1 - \cos\phi) & ac(1 - \cos\phi) \\ ba(1 - \cos\phi) & b^2 + (c^2 + a^2)\cos\phi & bc(1 - \cos\phi) \\ ca(1 - \cos\phi) & cb(1 - \cos\phi) & c^2 + (a^2 + b^2)\cos\phi \end{vmatrix}$

is independent of a, b, c

11.

Show that the value of the determinant $\begin{vmatrix} \tan(A+P) & \tan(B+P) & \tan(C+P) \\ \tan(A+Q) & \tan(B+Q) & \tan(C+Q) \\ \tan(A+R) & \tan(B+R) & \tan(C+R) \end{vmatrix}$ vanishes for all values of A, B, C, P, Q & R where $A + B + C + P + Q + R = 0$.

12. Prove that $\begin{vmatrix} -bc & b^2 + bc & c^2 + bc \\ a^2 + ac & -ac & c^2 + ac \\ a^2 + ab & b^2 + ab & -ab \end{vmatrix} = (ab + bc + ca)^3.$

13. Show that, $\begin{vmatrix} \cos(x+x^2) & \sin(x+x^2) & -\cos(x+x^2) \\ \sin(x-x^2) & \cos(x-x^2) & \sin(x-x^2) \\ \sin 2x & 0 & \sin 2x^2 \end{vmatrix} = \sin(2x+2x^2).$

14. If $\begin{vmatrix} \frac{1}{a+x} & \frac{1}{b+x} & \frac{1}{c+x} \\ \frac{1}{a+y} & \frac{1}{b+y} & \frac{1}{c+y} \\ \frac{1}{a+z} & \frac{1}{b+z} & \frac{1}{c+z} \end{vmatrix} = \frac{P}{Q}$ where Q is the product of the denominators, prove that

$$P = (a-b)(b-c)(c-a)(x-y)(y-z)(z-x)$$

If A_1, B_1, C_1, \dots are respectively the cofactors of the elements a_1, b_1, c_1, \dots of the determinant

$$\Delta = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

then prove that

(i) $\begin{vmatrix} B_2 & C_2 \\ B_3 & C_3 \end{vmatrix} = a_1 \Delta.$

(ii) $\begin{vmatrix} A_1 & B_1 & C_1 \\ A_2 & B_2 & C_2 \\ A_3 & B_3 & C_3 \end{vmatrix} = \Delta^2$

16. Show that, $\begin{vmatrix} bc - a^2 & ca - b^2 & ab - c^2 \\ ca - b^2 & ab - c^2 & bc - a^2 \\ ab - c^2 & bc - a^2 & ca - b^2 \end{vmatrix} = \begin{vmatrix} a^2 & c^2 & 2ac - b^2 \\ 2ab - c^2 & b^2 & 2bc - a^2 \\ b^2 & c^2 & a^2 \end{vmatrix}$

17. Using consistency of equations, prove that if $bc + qr = ca + rp = ab + pq = -1$ then $\begin{vmatrix} ap & a & p \\ bq & b & q \\ cr & c & r \end{vmatrix} = 0.$

18. Show that : $\begin{vmatrix} \sin \alpha & \cos \alpha & 1 \\ \sin \beta & \cos \beta & 1 \\ \sin \gamma & \cos \gamma & 1 \end{vmatrix} = \sin(\alpha - \beta) + \sin(\beta - \gamma) + \sin(\gamma - \alpha).$

19. If $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = (l_1x + m_1y + n_1)(l_2x + m_2y + n_2)$, then prove that

$$\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = 0.$$

20. Find all the values of t for which the system of equations;

$$\begin{aligned} (t-1)x + (3t+1)y + 2tz &= 0 \\ (t-1)x + (4t-2)y + (t+3)z &= 0 \\ 2x + (3t+1)y + 3(t-1)z &= 0 \end{aligned}$$

has non trivial solutions and in this context find the ratios of x: y: z, when t has the smallest of these values.

21. Let $a > 0, d > 0$. Find the value of determinant

$$\begin{vmatrix} \frac{1}{a} & \frac{1}{a(a+d)} & \frac{1}{(a+d)(a+2d)} \\ \frac{1}{(a+d)} & \frac{1}{(a+d)(a+2d)} & \frac{1}{(a+2d)(a+3d)} \\ \frac{1}{(a+2d)} & \frac{1}{(a+2d)(a+3d)} & \frac{1}{(a+3d)(a+4d)} \end{vmatrix}.$$

[IIT – 1996, 5]

22. Let a, b, c be real numbers with $a^2 + b^2 + c^2 = 1$. Show that the equation

$$\begin{vmatrix} ax - by - c & bx + ay & cx + a \\ bx + ay & -ax + by - c & cy + b \\ cx + a & cy + b & -ax - by + c \end{vmatrix} = 0$$

represents a straight line

[IIT – 2001, 6]

Exercise - 7

1. D 2. D 3. B 4. B 5. D 6. B 7. B
8. A 9. B 10. B 11. B 12. B 13. A 14. A
15. A 16. AB 17. AB 18. BD 19. ABD 20. ABD
21. BD

Exercise - 8

1. (i) 0 (ii) $5(3\sqrt{2} - 5\sqrt{3})$
2. $x = -2 b/a$
6. $a = -2$
7. (a) $\lambda \neq 3$ (b) $\lambda = 3, \mu = 10$ (c) $\lambda = 3, \mu \neq 10$
8. for $c = 0, x = -3, y = 3$; for $c = -10, x = -\frac{1}{2}, y = \frac{4}{3}$

20. $t = 0$ or $3; x: y: z = 1: 1: 1$

21.
$$\frac{4d^4}{a(a+d)^2(a+2d)^3(a+3d)^2(a+4d)}$$

Exercise - 9

Part : (A) Only one correct option

1. Let a, b, c, d, u, v be integers. If the system of equations $ax + by = u, cx + dy = v$ has a unique solution in integers, then
 (A) $ad - bc = 1$
 (C) $ad - bc \neq 0$
 If $AB = O$ for the matrices
2. $A = \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$ and $B = \begin{bmatrix} \cos^2 \phi & \cos \phi \sin \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix}$ then $\theta - \phi$ is
 (A) an odd multiple of $\frac{\pi}{2}$
 (B) an even multiple of $\frac{\pi}{2}$
 (C) an odd multiple of π
 (D) 0
3. If $X = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, then value of X^n is
 (A) $\begin{bmatrix} 3n & -4n \\ n & -n \end{bmatrix}$
 (B) $\begin{bmatrix} 2+n & 5-n \\ n & -n \end{bmatrix}$
 (C) $\begin{bmatrix} 3^n & (-4)^n \\ 1^n & (-1)^n \end{bmatrix}$
 (D) none of these
4. If the matrix $\begin{bmatrix} 0 & 2\beta & \gamma \\ \alpha & \beta & -\gamma \\ \alpha & -\beta & \gamma \end{bmatrix}$ is orthogonal, then
 (A) $\alpha = \pm \frac{1}{\sqrt{2}}$
 (B) $\beta = \pm \frac{1}{\sqrt{6}}$
 (C) $\gamma = \pm \frac{1}{\sqrt{3}}$
 (D) all of these
5. If A, B are two $n \times n$ non-singular matrices, then
 (A) AB is non-singular
 (C) $(AB)^{-1} = A^{-1} B^{-1}$
 If B is a non-singular matrix and A is a square matrix, then $\det(B^{-1}AB)$ is equal to
 (A) $\det(A^{-1})$
 (B) $\det(B^{-1})$
 (C) $\det(A)$
 (D) $\det(B)$
6. If A is a square matrix of order $n \times n$ and k is a scalar, then $\text{adj}(kA)$ is equal to
 (A) $k \text{adj } A$
 (B) $k^n \text{adj } A$
 (C) $k^{n-1} \text{adj } A$
 (D) $k^{n+1} \text{adj } A$
7. Let A be a matrix of rank r. Then
 (A) $\text{rank}(A^T) = r$
 (B) $\text{rank}(A^T) < r$
 (C) $\text{rank}(A^T) > r$
 (D) none of these
8. If A = dig (2, -1, 3), B = dig (-1, 3, 2), then $A^2B =$
 (A) dig (5, 4, 11)
 (B) dig (-4, 3, 18)
 (C) dig (3, 1, 8)
 (D) B
9. If ω is a cube root of unity and $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{bmatrix}$, then $A^{-1} =$

$$(A) \begin{bmatrix} 1 & \omega & \omega^2 \\ \omega^2 & 1 & \omega \\ \omega & \omega^2 & 1 \end{bmatrix} \quad (B) \frac{1}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{bmatrix} \quad (C) \begin{bmatrix} 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \\ 1 & 1 & 1 \end{bmatrix} \quad (D) \frac{1}{2} \begin{bmatrix} 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \\ 1 & 1 & 1 \end{bmatrix}$$

11. If the system of equations $ax + y + z = 0$, $x + by + z = 0$ and $x + y + cz = 0$, where

$a, b, c \neq 1$, has a non-trivial solution, then the value of $\frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c}$ is:

- (A) -1 (B) 0 (C) 1 (D) None of these

12. If A is a square matrix of order 3, then the true statement is (where I is unit matrix).

- (A) $\det(-A) = -\det A$ (B) $\det A = 0$
 (C) $\det(A + I) = 1 + \det A$ (D) $\det 2A = 2 \det A$

13. Which of the following is incorrect

- (A) $A^2 - B^2 = (A + B)(A - B)$ (B) $(A^T)^T = A$
 (C) $(AB)^n = A^n B^n$, where A, B commute (D) $(A - I)(I + A) = O \Leftrightarrow A^2 = I$

14. The value of a for which system of equations, $a^3x + (a+1)^3y + (a+2)^3z = 0$,

$ax + (a+1)y + (a+2)z = 0$, $x + y + z = 0$, has a non-zero solution is:

- (A) -1 (B) 0 (C) 1 (D) none of these

15. If $A = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix}$ then AB is equal to

- (A) B (B) $3B$ (C) B^3 (D) $A + B$

16. If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ satisfies the equation $x^2 - (a+d)x + k = 0$, then

- (A) $k = bc$ (B) $k = ad$
 (C) $k = a^2 + b^2 + c^2 + d^2$ (D) $ad - bc$

17. Let $A = \begin{bmatrix} x+\lambda & x & x \\ x & x+\lambda & x \\ x & x & x+\lambda \end{bmatrix}$, then A^{-1} exists if

- (A) $x \neq 0$ (B) $\lambda \neq 0$ (C) $3x + \lambda \neq 0, \lambda \neq 0$ (D) $x \neq 0, \lambda \neq 0$

18. Identify the correct statement
 (A) If system of n simultaneous linear equations has a unique solution, then coefficient matrix is singular
 (B) If system of n simultaneous linear equations has a unique solution, then coefficient matrix is non singular
 (C) If A^{-1} exists, $(\text{adj } A)^{-1}$ may or may not exist

(D) $F(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 0 \end{bmatrix}$, then $F(x) \cdot F(y) = F(x-y)$

19. Matrix $A = \begin{bmatrix} x & 3 & 2 \\ 1 & y & 4 \\ 2 & 2 & z \end{bmatrix}$, if $xyz = 60$ and $8x + 4y + 3z = 20$, then $A(\text{adj } A)$ is equal to

$$(A) \begin{bmatrix} 64 & 0 & 0 \\ 0 & 64 & 0 \\ 0 & 0 & 64 \end{bmatrix} \quad (B) \begin{bmatrix} 88 & 0 & 0 \\ 0 & 88 & 0 \\ 0 & 0 & 88 \end{bmatrix} \quad (C) \begin{bmatrix} 68 & 0 & 0 \\ 0 & 68 & 0 \\ 0 & 0 & 68 \end{bmatrix} \quad (D) \begin{bmatrix} 34 & 0 & 0 \\ 0 & 34 & 0 \\ 0 & 0 & 34 \end{bmatrix}$$

20. If $P = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$, $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $Q = PAP^{-1}$ and $x = P^1 Q^{2005} P$, then x is equal to [IIT JEE - 2005]

(A) $\begin{bmatrix} 1 & 2005 \\ 0 & 1 \end{bmatrix}$

(C) $\frac{1}{4} \begin{bmatrix} 2+\sqrt{3} & 1 \\ -1 & 2-\sqrt{3} \end{bmatrix}$

(B) $\begin{bmatrix} 4+2005\sqrt{3} & 6015 \\ 2005 & 4-2005\sqrt{3} \end{bmatrix}$

(D) $\frac{1}{4} \begin{bmatrix} 2005 & 2-\sqrt{3} \\ 2+\sqrt{3} & 2005 \end{bmatrix}$

[IIT JEE - 2006]

Comprehension

$A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 2 & 1 \end{bmatrix}$, if U_1 , U_2 , and U_3 are columns matrices satisfying $AU_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $AU_2 = \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}$ and

$AU_3 = \begin{bmatrix} 2 \\ 3 \\ 1 \end{bmatrix}$. If U is 3×3 matrix whose columns are U_1 , U_2 , U_3 then answer the following questions

21. The value of $|U|$ is (A) 3 (B) -3 (C) 3/2 (D) 2 [IIT JEE - 2006]
22. The sum of the elements of U^{-1} is (A) -1 (B) 0 (C) 1 (D) 3 [IIT JEE - 2006]
23. The value of $[3 \ 2 \ 0] U \begin{bmatrix} 3 \\ 2 \\ 0 \end{bmatrix}$ is (A) 5 (B) 5/2 (C) 4 (D) 3/2 [IIT JEE - 2006]

Part : (B) May have more than one options correct

24. The rank of the matrix $\begin{bmatrix} -1 & 2 & 5 \\ 2 & -4 & a-4 \\ 1 & -2 & a+1 \end{bmatrix}$ is (A) 2 if $a = 6$ (B) 2 if $a = 1$ (C) 1 if $a = 2$ (D) 1 if $a = -6$
25. Which of the following statement is always true
 (A) Adjoint of a symmetric matrix is a symmetric matrix
 (B) Adjoint of a unit matrix is unit matrix
 (C) $A(\text{adj } A) = (\text{adj } A)A$
 (D) Adjoint of a diagonal matrix is diagonal matrix

26. Matrix $\begin{bmatrix} a & b & (ab-b) \\ b & c & (bc-c) \\ 2 & 1 & 0 \end{bmatrix}$ is non invertible if (A) $a = 1/2$ (B) a, b, c are in A.P. (C) a, b, c are in G.P. (D) a, b, c are in H.P.
27. The singularity of matrix $\begin{bmatrix} 1 & a & a^2 \\ \cos(p-d)x & \cos px & \cos(p+d)x \\ \sin(p-d)x & \sin px & \sin(p+d)x \end{bmatrix}$ depends upon which of the following parameter (A) a (B) p (C) x (D) d
28. Which of the following statement is true
 (A) Every skew symmetric matrix of odd order is non singular
 (B) If determinant of a square matrix is nonzero, then it non singular
 (C) Rank of a matrix is equal or higher than the order of the matrix
 (D) Adjoint of a singular matrix is always singular

29. If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ (where $bc \neq 0$) satisfies the equations $x^2 + k = 0$, then
 (A) $a + d = 0$ (B) $k = -|A|$ (C) $k = |A|$ (D) none of these

30. If $A^{-1} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & -2 & 1 \\ 0 & 0 & -1 \end{bmatrix}$, then
 (A) $|A| = 2$ (B) A is non-singular
 (C) $\text{Adj. } A = \begin{bmatrix} 1/2 & -1/2 & 0 \\ 0 & -1 & 1/2 \\ 0 & 0 & -1/2 \end{bmatrix}$ (D) A is skew symmetric matrix

Exercise - 10

1. Find x so that $\begin{bmatrix} 1 & x & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 3 & 2 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 0$
2. If A and B are two square matrices such that $AB = A$ & $BA = B$, prove that A & B are idempotent
3. If $f(x) = x^2 - 5x + 7$, find $f(A)$ where $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$.
4. Prove that the product of matrices

Get Solution of These Packages & Learn by Video Tutorials on www.MathsBySuhag.com

$\begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$ and $\begin{bmatrix} \cos^2 \phi & \cos \phi \sin \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix}$ is the null matrix, when θ and ϕ differ by an odd multiple of $\pi/2$.

Given $F(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$. If $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$. Then for what values of y ,
 $F(x+y) = F(x)F(y)$.

Find the values of x, y, z if the matrix $A = \begin{bmatrix} 0 & 2y & z \\ x & y & -z \\ x & -y & z \end{bmatrix}$ obeys the law $A^t A = I$.

Compute A^{-1} for the following matrix $A = \begin{bmatrix} -1 & 2 & 5 \\ 2 & -3 & 1 \\ -1 & 1 & 1 \end{bmatrix}$. Hence solve the system of equations;
 $-x + 2y + 5z = 2; 2x - 3y + z = 15 \text{ & } -x + y + z = -3$

Show that $\begin{bmatrix} 1 & -\tan \theta/2 \\ \tan \theta/2 & 1 \end{bmatrix} \begin{bmatrix} 1 & \tan \theta/2 \\ -\tan \theta/2 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$

Gaurav purchases 3 pens, 2 bags and 1 instrument box and pays Rs. 41. From the same shop Dheeraj purchases 2 pens, 1 bag and 2 instrument boxes and pays Rs. 29, while Ankur purchases 2 pens, 2 bags and 2 instrument boxes and pays Rs. 44. Translate the problem into a system of equations. Solve the system of equations by matrix method and hence find the cost of 1 pen, 1 bag and 1 instrument box.

If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, then prove that $A^2 - 4A - 5I = 0$.

(a) using A^{-1} Having given equations $x = c_1 y + b_1 z, y = a_1 z + c_1 x, z = b_1 x + a_1 y$ where x, y, z are not all zero, prove that $a_1^2 + b_1^2 + c_1^2 + 2abc_1 - 1 = 0$.

Consider the system of linear equations in x, y, z :

$$\begin{aligned} (\sin 3\theta)x - y + z &= 0 \\ (\cos 2\theta)x + 4y + 3z &= 0 \\ 2x + 7y + 7z &= 0 \end{aligned}$$

Find the values of θ for which this system has non-trivial solution.

Solve the following systems of linear equations by using the principle of matrix.

(i) $\begin{aligned} 2x - y + 3z &= 8 \\ -x + 2y + z &= 4 \\ 3x + y - 4z &= 0 \end{aligned}$

(ii) $\begin{aligned} x + y + z &= 9 \\ 2x + 5y + 7z &= 52 \\ 2x + y - z &= 0 \end{aligned}$

Compute A^{-1} if $A = \begin{bmatrix} 3 & -2 & 3 \\ 2 & 1 & -1 \\ 4 & -3 & 2 \end{bmatrix}$. Hence solve the system of equations $\begin{bmatrix} 3 & 0 & 3 \\ 2 & 1 & 0 \\ 4 & 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 8 \\ 1 \\ 4 \end{bmatrix} + \begin{bmatrix} 2y \\ z \\ 3y \end{bmatrix}$.

Find the rank of the following matrices:

(i) $\begin{bmatrix} 1 & 2 & 3 & 2 \\ 2 & 3 & 5 & 1 \\ 1 & 3 & 4 & 5 \end{bmatrix}$

(ii) $\begin{bmatrix} 1 & 2 & -1 & 3 \\ 4 & 1 & 2 & 1 \\ 3 & -1 & 1 & 2 \\ 1 & 2 & 0 & 1 \end{bmatrix}$

(iii) $\begin{bmatrix} 1 & 3 & 4 & 3 \\ 3 & 9 & 12 & 3 \\ 1 & 3 & 4 & 1 \end{bmatrix}$

(iv) $\begin{bmatrix} 0 & 1 & -3 & -1 \\ 1 & 0 & 1 & 1 \\ 3 & 1 & 0 & 2 \\ 1 & 1 & -2 & 0 \end{bmatrix}$

Determine the product $\begin{bmatrix} -4 & 4 & 4 \\ -7 & 1 & 3 \\ 5 & -3 & -1 \end{bmatrix} \begin{bmatrix} 1 & -1 & 1 \\ 1 & -2 & -2 \\ 2 & 1 & 3 \end{bmatrix}$ and use it to solve the system of equations.

$x - y + z = 4; x - 2y - 2z = 9; 2x + y + 3z = 1$.

If $A = \begin{pmatrix} a & b & c \\ b & c & a \\ c & a & b \end{pmatrix}$, where a, b, c are real positive numbers, $abc = 1$ and $A^T A = I$, then find the value of $a^3 + b^3 + c^3$.

[IIT JEE - 2003, 2]

If M is 3×3 matrix M has its $\det(M) = 1$ and $MM^T = I$. Prove that $\det(M - I) = 0$.

[IIT JEE - 2004, 2]

19. If $A = \begin{bmatrix} a & 1 & 0 \\ 1 & b & d \\ 1 & b & c \end{bmatrix}$, $B = \begin{bmatrix} a & 1 & 1 \\ 0 & d & c \\ f & g & h \end{bmatrix}$, $U = \begin{bmatrix} f \\ g \\ h \end{bmatrix}$, $V = \begin{bmatrix} a^2 \\ 0 \\ 0 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$

and $AX = U$ has infinitely many solution. Prove that $BX = V$ has no unique solution, also prove that if $adf \neq 0$, then $BX = V$ has no solution.

[IIT JEE - 2004, 4]

Exercise - 9

1. C 2. A 3. D 4. D 5. A 6. C 7. C
8. A 9. B 10. B 11. C 12. A 13. A 14. A
15. B 16. D 17. C 18. B 19. C 20. A 21. A
22. B 23. A 24. ABD 25. ABCD 26. AB
27. CD 28. BD 29. AC 30. BC

7. $A^{-1} = -\frac{1}{7} \begin{bmatrix} -4 & 3 & 17 \\ -3 & 4 & 11 \\ -1 & -1 & -1 \end{bmatrix}$ & $x = 2, y = -3, z = 2$

9. Rs. 2, Rs. 15 & Rs. 5

12. $\theta = n\pi, n\pi + (-1)^n \frac{\pi}{6}; n \in \mathbb{Z}$

13. (i) $x = 2; y = 2; z = 2$ (ii) $x = 1; y = 3; z = 5$

14. $x = 1; y = 2; z = 3$

15. (i) 2 (ii) 3 (iii) 2 (iv) 2

16. $x = 3; y = -2; z = -1$

17. 4

Exercise - 10

1. $-\frac{9}{8}$
3. $f(A) = 0$
5. $y \in \mathbb{R}$

6. $x = \pm \frac{1}{\sqrt{2}}, y = \pm \frac{1}{\sqrt{6}}, z = \pm \frac{1}{\sqrt{3}}$